FORM NHAMCS-906 (9-14-2009) U.S. DEPARTMENT OF COMMERCE
Economics and Statistics Administration
U.S. CENSUS BUREAU
ACTING AS DATA COLLECTION AGENT FOR THE
U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

National Center for Health Statistics

NATIONAL HOSPITAL AMBULATORY MEDICAL CARE SURVEY 2010 CERVICAL CANCER SCREENING SUPPLEMENT

NOTICE – Public reporting burden of this collection of information is estimated to average 15 minutes per response, including the time f or reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently v alid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office; 1600 Clifton Road, MS D-74, Atlanta, GA 30333, ATTN: PRA (0920-0234).

Assurance of Confidentiality – All information which would permit identification of an individual, a practice, or an establishment will be held confidential, will be used for statistical purposes only by NCHS staff, contractors, and agents only when required and with necessary controls, an divil not be disclosed or released to other persons without the consent of the individual or establishment in accordance with section 308(d) of the Public Health S ervice Act (42 USC 242m) and the Confidential Information Protection and Statistical Efficiency Act (PL-107-347).

	BACKGROUND	INFORMATION								
A. Hospital number	only ONE.	C. AU number								
D. Census contact name	E. Census conta telephone	act	Area code Number							
The Centers for Disease Control and Prevention is conducting a special survey on cervical cancer screening performed in hospital outpatient clinics. Please answer the following questions. We appreciate your time on this important public health concern.										
1. Does this clinic use any of the fol	lowing methods to screen for		Mark (X) one interval for routine screening.							
cervical cancer? Mark (X) all that apply. a. Conventional Pap test (Definition		 Annually 	Every 2 years	Every 3 years	More than 3 years	No routine interval recom- mended				
1 Yes – How often does this c method? — 2 No 3 Unknown Continue with it		1	2	3 🗆	4	5				
b. Liquid-based cytology (Definition 1 Yes – How often does this c method? 2 No 3 Unknown Continue with it		 	2	3□	4	5				
c. Other – <i>Specify</i>	c. Other – Specify ⊋									
1 ☐ Yes – How often does this c method? — 2 ☐ No 3 ☐ Unknown	linic routinely screen women	using this	 1	2	3□	4	5			
2. Does this clinic perform colposed 1 Yes 2 No 3 Unknown	opy?									

	Does this clinic ever order or collect the Human Papillomavirus (HPV) DNA test? 1 Yes - Go to item 3b 2 No - SKIP to item 3c 3 Not aware of HPV DNA test 4 Unknown Which of the following HPV DNA tests are ordered or collected in this clinic? Mark (X) all that apply. 1 High risk (HR) HPV DNA test 2 Low risk (LR) HPV DNA test	4a.	If a patient's Pap test result is borderline or abnormal, does this clinic routinely order an HPV DNA test to be performed on that sample (commonly called reflex HPV DNA testing)? (An HPV DNA test may be run on the same liquid-based medium as the Pap test or an HPV DNA test specimen may be collected at the same time as the conventional Pap test.) 1 Yes - Go to item 4b 2 No 3 Unknown 3 SKIP to item 5a				
	SKIP to low risk HPV DNA test 3 □ Not aware there was a high risk or low risk HPV DNA test 4 □ Type-specific HPV DNA test 5 □ Unknown	b.	For which borderline or abnormal Pap test result would this clinic order or collect a reflex HPV DNA test? Mark (X) all that apply. 1 ASC-US (atypical squamous cells of undetermined significance)				
c.	 Why is the HPV DNA test not ordered or collected in this clinic? – <i>Mark (X) all that apply.</i> □ This clinic does not see the types of patients for whom the HPV DNA test is indicated. □ This clinic uses other tests, procedures, or examination methods to manage patients for whom the HPV DNA test is indicated. □ The patients in this clinic have timely access to 		 ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion) LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1) HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ) AGC (atypical glandular cells) 				
	colposcopy. 4 Assessing patients' HPV infection status is not a priority at this clinic. 5 The labs affiliated with this clinic do not offer the HPV DNA test. 6 The health plans or health systems affiliated with this clinic do not recommend the HPV DNA test. 7 The HPV DNA test is not a reimbursed or covered	c.	For which patients does this clinic usually order reflex HPV DNA testing? — Mark (X) all that apply. 1 ☐ Women under 21 years old 2 ☐ Women 21 years old to 29 years old 3 ☐ Women 30 years old and over 4 ☐ Other — Specify ✓				
	service for most patients in this clinic. 8 Discussing cervical cancer screening in the context of an STD is avoided in this clinic. 9 Notifying or counseling patients about positive HPV DNA test results would take too much time. 10 Notifying or counseling patients about positive HPV DNA test results might make clinicians in this clinic	5a.	Does this clinic routinely recall patients to come back for a second sample collection for an HPV DNA test if their Pap test is abnormal or borderline (recall testing)? 1 Yes – Go to item 5b 2 No 3 Unknown SKIP to item 6a on page 3				
	feel uncomfortable. 11 Notifying or counseling patients about positive HPV DNA test results might make patients in this clinic feel uncomfortable, angry, or upset. SKIP to item 7 on page 3.		For which abnormal or borderline Pap test result would this clinic recall a patient for an HPV DNA test? Mark (X) all that apply. 1 ASC-US (atypical squamous cells of undetermined significance) 2 ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion) 3 LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1) 4 HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ)				
			5 AGC (atypical glandular cells)				

Page 2 FORM NHAMCS-906 (9-14-2009)

6a.	6a. Does this clinic routinely order or collect an HPV DNA test at the same time as the Pap test as part of routine cervical cancer screening (commonly called adjunct HPV testing or cotesting)?										
	1 ☐ Yes – Go to item 6b										
	2 No CKID to item 7										
	3 Unknown SKIP to item 7										
b.	For which patients does this clinic routinely order or collect an HPV DNA test along with the Pap test (commonly called adjunct HPV testing or cotesting)? Mark (X) all that apply.									ed	
	1 🗆 '	Women under 21 yea	ars old								
		Women 21 years old		s old							
	3 🗌	Women 30 years old	and over								
		Women who request									
		Women who request	the test to	check the	ir HPV inted	tion status					
	6 🗀 '	Other – Specify 📈									
7.	Giv	ven the following scre	enina histo	ries, wher	n would this	clinic recom	mend that a	woman be	tween 30 au	nd 60	
	yea	ars of age return for h	ner next Pa	p test?							
			 		For eac	ch of the fol	lowing scena	arios, mark	(X) only C	NE for each	row.
Pr	ior Pa	ap test results Ist 5 years	Current	Current							Have no experience
	(excl	uding current	DNA test	Pap test	No follow-up	Less than	6 months to less	1	2	3 years	with this
	nori	mal results)	results	result	needed	6 months	than 1 year	year	years	or more	type of patient or
			 								test
	(a)	Two consecutive normal Pap tests	Has not had test	Normal	1 🗆	2	3 🗆	4	5 🗆	6	7
	<u></u>	Two consecutive									,
	(0)	normal Pap tests	Negative	Normal	1 🗌	2	3 🗌	4	5 🗌	6	7
	(c)	Two consecutive normal Pap tests	 	Nissensel	1 🗆	2	3 🗆	4	5 🔲	6	7 🗆
			Positive	Normal	''		30		3		, ,
	(d)	Has not had a Pap test	 Negative	Normal	1 🗆	2	3□	4	5 🗆	6□	7 🗆
	_		· · · · · · · · · · · · · · · · · · ·								
	(e)	Has not had a Pap	Positive	Normal	1 🗆	2	3□	4	5□	6 🗆	7 🗆
		test	1 0311110	Nomai							
	(f)	Abnormal Pap test	Negative	Normal	1 🗆	2	3□	4	5 🗆	6□	7 🗆
	(g)	Abnormal Pap test	Positive	Normal	1 🗌	2	3 🗌	4	5 🗌	6 🗆	7 🗌
QUESTIONS 8-14 ASK ABOUT THE HPV VACCINE											
8.	8. How often does this clinic use an HPV test to determine who should get the HPV vaccine? Mark (X) only one.										
	1 ☐ Rarely or never										
		Sometimes									
		Usually	lwovo								
		Always or almost a Do not recommend	-	accine – S	KIP to item	10.					
	5 ☐ Do not recommend the HPV vaccine – SKIP to item 10.										

FORM NHAMCS-906 (9-14-2009) Page 3

9.	As it relates to the HPV vaccine,								
	how often does your clinic –	Rarely or	Sometimes	Usually	Always or almost	Unknown/Not applicable/ Do not ask			
	Mark (X) only ONE for each row.	never		,	always				
	a. Use the number of sexual partners to determine who should get the HPV vaccine?	1 🗆	2	3□	4 🗆	5 🗌			
	b. Perform a Pap test to determine who should get the HPV vaccine?	1	2	3 🗆	4 🗆	5 🔲			
	c. Recommend the HPV vaccine to females with a history of an abnormal Pap test result (ASC-US or higher)?	1 🗆	2 🗆	3□	4 🗆	5 🗌			
	d. Recommend the HPV vaccine to females with a positive HPV test?	1 🗌	2	3 🗆	4 🗆	5 🗌			
10.	Mill this clinic's cervical cancer screening and management procedures change for females who have been fully vaccinated with the HPV vaccine? 1 Yes 2 No − SKIP to item 14								
11.	How will this clinic determine when to start ro cervical cancer screening for fully HPV vaccifemales?	age as non-HPV ed females –							
	Mark (X) all that apply.	1	Specify		-				
		i	2∐ At a later Specify		→				
			2 By onset of s	sexual activity – ear(s) since ual activity?———>					
		İ	onset of sexu						
		ļ	3 Will not be so 4 Unknown	reening fully HPV vaccinated females					
12.	How often will this clinic routinely screen for	often will this clinic routinely screen for cervical							
	cancer among females that have been fully vaccinated with the HPV vaccine? Mark (X) one. 2 Every 2–3 years 3 Every 4–5 years								
	, ,	4□ Greater than every 5 years							
13.	■ Will this clinic be using the HPV DNA test for managing abnormal cytology for females that have been fully vaccinated with the HPV vaccine?								
14.	Please indicate whether you agree with, disa unsure of the statements in a. and b.	Agree	Disagree	Unsure					
	a. There will be fewer numbers of abnormation among vaccinated females.	1 🗆	2	3□					
	b. There will be fewer referrals for colpose vaccinated females.	1 🗆	2	3□					
15.	The Centers for Disease Control and Prevention (CDC) funds state health departments to provide breast and cervical cancer screening services to low income women through the National Breast and Cervical Cancer Early Detection Program (Title XV). The state health departments contract out the screening services to physicians and other health care providers. Is this clinic currently participating in this state or national screening program?								
	1 Yes 2 No 3 Unknown								
16.	 For purposes of this survey, which of the following categories describe your profession? – Mark (X) only ONE. 1 Physician 2 Physician assistant/ 3 Registered nurse 4 Other clinic staff Nurse practitioner/ Nurse midwife 								

CLOSING STATEMENT

Thank you for completing this special survey. We appreciate your time and cooperation.

Page 4 FORM NHAMCS-906 (9-14-2009)