FORM NHAMCS-906

U.S. DEPARTMENT OF COMMERCE Economics and Statistics Administration U.S. CENSUS BUREAU ACTING AS DATA COLLECTION AGENT FOR THE U.S. Department of Health and Human Services Centers for Disease Control and Prevention National Center for Health Statistics

NATIONAL HOSPITAL AMBULATORY MEDICAL CARE SURVEY 2008 CERVICAL CANCER SCREENING SUPPLEMENT

NOTICE – Public reporting burden of this collection of information is estimated to average 15 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to: CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road, MS D-74, Atlanta, GA 30333, ATTN: PRA(0920-0234).

Assurance of Confidentiality - All information which would permit identification of an individual, a practice, or an establishment will be held confidential.

will be u without t	sed by persons engaged in and for the the consent of the individual or the esta	purpose of the survey and will not blishment in accordance with section	be disclosed or on 308(d) of the	released Public He	to other per ealth Service	sons or use Act (42 US	d for any oth C 242m).	ner purpose			
		BACKGROUND	INFORMA	TION							
0010	- Hospital number	ODIS B. Clinic type – Ma			0020 C.	AU nun	nber				
0025 D	D. Census contact name 0030 E. Census telepho						contact Area code Number				
INTR	SCREENING P	rs for Disease Control and performed in hospital outpa your time on this importan	atient clinics.	Please	e answer ern.	the follow	ving ques	stions. W	e		
(1. D	oes this clinic use any of the fo	llowing methods to screen f	for		Mark (X	lark (X) one interval for routine screening.					
cervical cancer? Mark (X) all that apply. a. Conventional Pap test (Definition – Smear spread on glass slide and fixed)						Every 2 years	Every 3 years	More than 3 years	No routine interval recom- mended		
1 Yes – How often does this clinic routinely screen women using this method? 2 No 3 Unknown Continue with item 1b					1 1 1 1 1 1 1 1 1 1 1	2	3	4	5 🗌		
b. Liquid-based cytology (Definition – Specimen suspended in liquid solution) Ooe						2	3	4	5		
c. Ot	ther – <i>Specify</i>				l I						
5005		linia routinaly caroon waman	using this		 						
2	1 Yes - How often does this clinic routinely screen women using this method? 2 No 3 Unknown					2	3 🗆	4	5 🗌		
0050 1	Ooes this clinic perform colposod ☐ Yes ☐ No ☐ Unknown	opy?									

0070 b.	Does this clinic ever order or collect the Human Papillomavirus (HPV) DNA test? 1 Yes - Go to item 3b 2 No - SKIP to item 3c 3 Not aware of HPV DNA test 4 Unknown Which of the following HPV DNA tests are ordered or collected in this clinic? Mark (X) all that apply. 1 High risk (HR) HPV DNA test 2 Low risk (LR) HPV DNA test	0085	If a patient's Pap test result is borderline or abnormal, does this clinic routinely order an HPV DNA test to be performed on that sample (commonly called reflex HPV DNA testing)? (An HPV DNA test may be run on the same liquid-based medium as the Pap test or an HPV DNA test specimen may be collected at the same time as the conventional Pap test.) 1 Yes – Go to item 4b 2 No 3 Unknown 3 KIP to item 5a				
	3 ☐ Not aware there was a high risk or low risk HPV DNA test 4 ☐ Type-specific HPV DNA test 5 ☐ Unknown	b.	For which borderline or abnormal Pap test result would this clinic order or collect a reflex HPV DNA test? Mark (X) all that apply. 1 ASC-US (atypical squamous cells of undetermined significance)				
0080	Why is the HPV DNA test not ordered or collected in this clinic? − <i>Mark</i> (<i>X</i>) all that apply. 1 This clinic does not see the types of patients for whom the HPV DNA test is indicated. 2 This clinic uses other tests, procedures, or examination methods to manage patients for whom the HPV DNA test is indicated. 3 The patients in this clinic have timely access to		2 ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion) 3 LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1) 4 HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ) 5 AGC (atypical glandular cells)				
	colposcopy. 4 ☐ Assessing patients' HPV infection status is not a priority at this clinic. 5 ☐ The labs affiliated with this clinic do not offer the HPV DNA test. 6 ☐ The health plans or health systems affiliated with	0095	For which patients does this clinic usually order reflex HPV DNA testing? – Mark (X) all that apply. 1 Women under 21 years old 2 Women 21 years old to 29 years old 3 Women 30 years old and over 4 Other – Specify				
	this clinic do not recommend the HPV DNA test. 7 The HPV DNA test is not a reimbursed or covered service for most patients in this clinic.	5010					
	8 ☐ Discussing cervical cancer screening in the context of an STD is avoided in this clinic.	5a.	Does this clinic routinely recall patients to come back for a second sample collection for an HPV DNA test if their Pap test is abnormal or borderline (recall testing)?				
	 Notifying or counseling patients about positive HPV DNA test results would take too much time. Notifying or counseling patients about positive HPV DNA test results might make clinicians in this clinic feel uncomfortable. Notifying or counseling patients about positive HPV DNA test results might make patients in this clinic feel uncomfortable, angry, or upset. 	b.	1 ☐ Yes - Go to item 5b 2 ☐ No 3 ☐ Unknown } SKIP to item 6a on page 3 For which abnormal or borderline Pap test result would this clinic recall a patient for an HPV DNA test? Mark (X) all that apply. 1 ☐ ASC-US (atypical squamous cells of undetermined significance)				
	SKIP to item 7 on page 3.		2 ☐ ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion) 3 ☐ LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1) 4 ☐ HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ) 5 ☐ AGC (atypical glandular cells)				

Page 2 FORM NHAMCS-906 (11-28-2007)

6a.	6a. Does this clinic routinely order or collect an HPV DNA test at the same time as the Pap test as part of routine cervical cancer screening (commonly called adjunct HPV testing or cotesting)?										
0110	1 ☐ Yes – Go to item 6b										
	2 No 3 Unknown SKIP to item 7										
b.	For which patients does this clinic routinely order or collect an HPV DNA test along with the Pap test (commonly called adjunct HPV testing or cotesting)? <i>Mark (X) all that apply.</i>										
0115											
	2 Women 21 years old to 29 years old										
	3 Women 30 years old and over										
	4 ☐ Women who request the test for cervical cancer screening 5 ☐ Women who request the test to check their HPV infection status										
		Other – <i>Specify</i>		000		0					
		ŕ									
5015											
7.	7. Given the following screening histories, when would this clinic recommend that a woman between 30 and 60 years of age return for her next Pap test?										
			 		For ea	ch of the fo	llowing scen	arios, mai	rk (X) only	ONE for ea	ch row.
i	n pa excl	ap test results est 5 years uding current mal results)	Current HPV DNA test results	Current Pap test result	No follow-up needed	Less than 6 months	6 months to less than 1 year	1 year	2 years	3 years or more	Have no experience with this type of patient or test
0120	(a)	Two consecutive normal Pap tests	Has not had test	Normal	1 🗆	2	3 🗆	4 🗌	5 🗆	6 🗆	7
0125	(b)	Two consecutive normal Pap tests	 Negative	Normal	1	2	3 🗆	4 🗌	5	6	7
0130	(c)	Two consecutive normal Pap tests	 Positive	Normal	1 🗆	2	з 🗆	4 🗌	5 🗆	6	7
0135	(d)	Has not had a Pap test	 Negative	Normal	1 🗌	2	з□	4	5	6	7
0140	(e)	Has not had a Pap test	Positive	Normal	1 🗆	2	з 🗆	4	5	6	7
0145	(f)	Abnormal Pap test	 Negative 	Normal	1 🗌	2	3 🗆	4	5	6	7 🗌
0150	(g)	Abnormal Pap test	Positive	Normal	1	2	3 🗆	4	5 🗌	6	7 🗌
QUESTIONS 8–14 ASK ABOUT THE HPV VACCINE											
8.	8. How often does this clinic use an HPV test to determine who should get the HPV vaccine? Mark (X) only one.										
	1 ☐ Rarely or never										
	2 Sometimes										
	3 ☐ Usually 4 ☐ Always or almost always										
	5 □ Do not recommend the HPV vaccine –SKIP to item 10.										

FORM NHAMCS-906 (11-28-2007) Page 3

9.	As it relates to the HPV vaccine.									
	how often does your clinic -	Rarely or never	Sometimes		Usually	Always or almost always	Unknown/Not applicable/			
	Mark (X) only ONE for each row.	lievei				aiways	Do not ask			
0120	a. Use the number of sexual partners to determine who should get the HPV vaccine?	1 🗌		2	з 🗆	4 🗆	5 🗌			
0125	b. Perform a Pap test to determine who should get the HPV vaccine?	1 🗆		2	3 🗆	4 🗆	5 🗌			
0130	c. Recommend the HPV vaccine to females with a history of an abnormal Pap test result (ASC-US or higher)?	1 🗆		2	3□	4	5 🗌			
0135	d. Recommend the HPV vaccine to females with a positive HPV test?	1 🗆		2	3□	4 🗆	5 🗌			
10.	O. Will this clinic's cervical cancer screening and management procedures change for females who have been fully vaccinated with the HPV vaccine? □ Yes □ No − SKIP to item 14									
11. How will this clinic determine when to start routine cervical cancer screening for fully HPV vaccinated females? 1 □ By age 1 □ At same age a vaccinated fer Nark (X) all that apply						→				
	Mark (X) all that apply.			Ороспу	.					
				2 ☐ At a later Specify						
		i	. 2		exual activity –					
		ļ		How many ye	year(s) since					
			onset of sexual activity?——→ ₃□ Will not be screening fully HPV vaccinated females							
		4 Unknown								
12.	How often will this clinic routinely screen for	cervical	1 1	Annually						
	cancer among females that have been fully vaccinated 2 Every 2-3 years									
	with the HPV vaccine? <i>Mark (X) one.</i>									
	l 4□ Greater than every 5 years 5□ Will not be screening fully HPV vaccinated females									
	6 Unknown									
13.	3. Will this clinic be using the HPV DNA test for managing abnormal cytology for females that have been fully vaccinated with the HPV vaccine? 1									
14.	Please indicate to what extent you agree, diswith each statement. Please respond to both	е	Agree	Disagree	Unsure					
	a. There will be fewer numbers of abnormal Pap tests among vaccinated females.				1 🗆	2	з 🗌			
	There will be fewer referrals for colpose vaccinated females.		1 🗆	2	3 🗆					
15.	15. The Centers for Disease Control and Prevention (CDC) funds state health departments to provide breast and cervical cancer screening services to low income women through the National Breast and Cervical Cancer Early Detection Program (Title XV). The state health departments contract out the screening services to physicians and other health care providers. Is this clinic currently participating in this state or national screening program?									
0155	155 1 Yes 2 No 3 Unknown									
16.	For purposes of this survey, which of the fo	llowing catego	ories d	escribe your p	orofession? - Ma	rk (X) only ONE.				
0160	1 □ Physician 2 □ Physician assistant/ 3 □ Registered nurse 4 □ Other clinic staff Nurse practitioner/ Nurse midwife									

CLOSING STATEMENT

Thank you for completing this special survey. We appreciate your time and cooperation.

Page 4 FORM NHAMCS-906 (11-28-2007)