


**Figure 1.1. Percentage of persons of all ages without health insurance coverage: United States, 1997–2005**



NOTES: A person was defined as uninsured if he or she did not have any private health insurance, Medicare, Medicaid, State Children's Health Insurance Program (SCHIP), State-sponsored or other government-sponsored health plan, or military plan at the time of the interview. A person was also defined as uninsured if he or she had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care. The analyses excluded persons with unknown health insurance status (about 1% of respondents each year). The data on health insurance status were edited using an automated system based on logic checks and keyword searches. For comparability, the estimates for all years were created using these same procedures. The resulting estimates of persons without health insurance coverage are generally 0.1–0.2 percentage points lower than those based on the editing procedures used for the final data files. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table II in the Appendix for more details.

In quarter 3 of 2004, two additional questions were added to the NHIS insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1 in the data table) and with the responses to these new questions (presented in the figure above and noted as Method 2 in the data table). Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, the percentage of uninsured persons was 14.4% (95% confidence interval = 13.5%–15.3%), which was not significantly lower than the Method 2 2004 estimate of 14.6%.
- The annual percentage of uninsured persons was 15.4% in 1997, 14.6% in 1998, 14.2% in 1999, and 14.9% in 2000. The estimate increased from 14.3% in 2001 to 15.2% in 2003, and then decreased (although not significantly) to 14.7% (Method 1) in 2004.

**Table 1.1. Number and percentage of persons without health insurance coverage, by age group: United States, 1997–2005**

Year	All ages	Under 65 years	18–64 years	Under 18 years	All ages	Under 65 years		18–64 years	Under 18 years
						Crude	Age-adjusted		
Number in millions					Percent (95% confidence interval)				
1997	410	40.7	30.8	9.9	15.4 (15.0-15.8)	17.4 (16.9-17.9)	17.2 (16.8-17.7)	18.9 (18.4-19.4)	13.9 (13.2-14.6)
1998	39.3	39.0	30.0	9.1	14.6 (14.1-15.1)	16.5 (16.0-17.0)	16.4 (15.9-16.9)	18.2 (17.7-18.7)	12.7 (12.0-13.4)
1999	38.7	38.3	29.8	8.5	14.2 (13.8-14.6)	16.0 (15.5-16.5)	16.0 (15.5-16.5)	17.8 (17.3-18.3)	11.8 (11.2-12.4)
2000	413	40.8	32.0	8.9	14.9 (14.5-15.3)	16.8 (16.3-17.2)	16.8 (16.3-17.3)	18.7 (18.1-19.2)	12.3 (11.7-12.9)
2001	402	39.8	31.9	7.9	14.3 (13.8-14.8)	16.2 (15.7-16.7)	16.2 (15.7-16.7)	18.3 (17.8-18.8)	11.0 (10.3-11.7)
2002	415	411	33.5	7.6	14.7 (14.3-15.1)	16.5 (16.0-16.9)	16.6 (16.1-17.1)	19.1 (18.6-19.6)	10.5 (9.9-11.1)
2003	436	43.2	35.9	7.3	15.2 (14.8-15.7)	17.2 (16.6-17.7)	17.3 (16.8-17.8)	20.1 (19.5-20.6)	10.1 (9.4-10.7)
2004 (Method 1 <sup>1</sup> )	42.5	42.0	35.0	7.0	14.7 (14.3-15.2)	16.6 (16.1-17.0)	16.7 (16.3-17.2)	19.4 (18.9-19.9)	9.6 (9.0-10.2)
2004 (Method 2 <sup>1</sup> )	42.1	41.7	34.9	6.8	14.6 (14.2-15.0)	16.4 (16.0-16.9)	16.6 (16.2-17.1)	19.3 (18.8-19.8)	9.4 (8.8-10.0)
Jan.–March 2005	417	41.4	35.1	6.3	14.4 (13.5-15.3)	16.2 (15.2-17.3)	16.4 (15.4-17.4)	19.3 (18.2-20.5)	8.6 (7.5-9.8)

<sup>1</sup> In quarter 3 of 2004, two additional questions were added to the NHIS insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1) and with the responses to these new questions (noted as Method 2). Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

NOTES: A person was defined as uninsured if he or she did not have any private health insurance, Medicare, Medicaid, State Children's Health Insurance Program (SCHIP), State-sponsored or other government-sponsored health plan, or military plan at the time of the interview. A person was also defined as uninsured if he or she had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care. The analyses excluded persons with unknown health insurance status (about 1% of respondents each year). The data on health insurance status were edited using an automated system based on logic checks and keyword searches. For comparability, the estimates for all years were created using these same procedures. The resulting estimates of persons without health insurance coverage are generally 0.1–0.2 percentage points lower than those based on the editing procedures used for the final data files. The number of uninsured persons was calculated as the percentage of uninsured persons multiplied by the total weighted population, including persons with unknown coverage. The age-specific numbers of uninsured may not add to their respective totals due to rounding error. Age-adjusted estimates for persons under 65 years of age for this Healthy People 2010 Leading Health Indicator are adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–44 years, and 45–64 years. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table II in the Appendix for more details.

DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, the percentage of uninsured persons was 16.2% (41.4 million) for persons under age 65 years, 19.3% (35.1 million) for persons aged 18–64 years, and 8.6% (6.3 million) for children under age 18 years.
- For adults aged 18–64 years, there was a decrease in the percentage of those who were uninsured from 1997 to 1999, followed by an increase from 1999 to 2003.

■ For children under age 18 years, the percentage of those who were uninsured decreased from 13.9% in 1997 to 8.6% in early 2005. This decreasing trend was observed regardless of whether Method 1 or Method 2 was used to calculate the 2004 estimate, but the extent of the decrease is larger using one method compared to the other method.

**Table 1.2. Percentage of persons under age 65 years with public health plan coverage and private health insurance coverage, by age group: United States, 1997–2005**

Year	Public <sup>1</sup>			Private <sup>1</sup>		
	Under 65 years	18–64 years	Under 18 years	Under 65 years	18–64 years	Under 18 years
	Percent (95% confidence interval)					
1997	13.6 (13.1-14.1)	10.2 (9.8-10.6)	21.5 (20.5-22.4)	70.8 (70.1-71.5)	72.8 (72.2-73.4)	66.2 (65.1-67.3)
1998	12.7 (12.2-13.2)	9.5 (9.1-9.9)	20.0 (19.0-20.9)	72.0 (71.3-72.7)	73.5 (72.9-74.1)	68.5 (67.4-69.5)
1999	12.4 (12.0-12.9)	9.0 (8.6-9.3)	20.5 (19.5-21.4)	73.1 (72.3-73.8)	74.8 (74.1-75.4)	69.1 (68.0-70.2)
2000	12.9 (12.4-13.4)	9.1 (8.7-9.4)	22.0 (21.0-23.0)	71.8 (71.1-72.5)	73.8 (73.2-74.4)	67.1 (66.1-68.2)
2001	13.6 (13.1-14.1)	9.4 (9.0-9.8)	23.6 (22.6-24.5)	71.6 (70.9-72.3)	73.7 (73.1-74.4)	66.7 (66.4-68.6)
2002	15.2 (14.6-15.8)	10.3 (9.9-10.7)	27.1 (26.0-28.2)	69.8 (69.0-70.6)	72.3 (71.6-72.9)	63.9 (62.7-65.1)
2003	16.0 (15.4-16.6)	10.9 (10.4-11.4)	28.6 (27.4-29.7)	68.2 (67.5-69.0)	70.6 (69.9-71.3)	62.6 (61.4-63.8)
2004 (Method 1 <sup>2</sup> )	16.1 (15.6-16.7)	11.1 (10.6-11.5)	28.5 (27.5-29.6)	68.6 (67.9-69.4)	70.9 (70.2-71.6)	63.1 (61.9-64.3)
2004 (Method 2 <sup>2</sup> )	16.2 (15.7-16.8)	11.1 (10.7-11.6)	28.7 (27.7-29.8)	...	...	...
Jan.–March 2005	17.1 (15.9-18.3)	11.5 (10.6-12.4)	30.8 (28.4-33.3)	68.0 (66.2-69.7)	70.5 (68.9-72.0)	61.8 (59.1-64.5)

...Category not applicable. See footnote 2 for more information.

<sup>1</sup>The category "public health plan coverage" includes Medicare (disability), Medicaid, State Children's Health Insurance Program (SCHIP), State-sponsored or other government-sponsored health plan, and military plans. The category "private health insurance" excludes plans that paid for only one type of service such as accidents or dental care. A small number of persons were covered by both public and private plans and were included in both categories. The analyses excluded persons with unknown health insurance status (about 1% of respondents each year). The data on type of coverage were edited using an automated system based on logic checks and keyword searches. For comparability, the estimates for all years were created using these same procedures. The resulting estimates of persons having public or private coverage are within 0.1–0.2 percentage points of those based on the editing procedures used for the final data files.

<sup>2</sup>In quarter 3 of 2004, two additional questions were added to the NHIS insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1) and with the responses to these new questions (noted as Method 2). Estimates of private insurance are not affected by the two additional questions. Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table II in the Appendix for more details.


DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, 17.1% of persons under age 65 years were covered by public plans, and 68.0% were covered by private health insurance plans.

■ For persons aged 18–64 years, after a period of decrease from 1997 to 1999, there was an increasing trend in public coverage from 1999 to early 2005. There was an increase from 1997 to 1999 followed by a decreasing trend from 1999 to early 2005 in private coverage for this age group. These trends were observed regardless of whether Method 1 or Method 2 was used to calculate the 2004 estimate, but the extent of the increase or decrease is larger using one method compared to the other method.

■ For children under age 18 years, after a period of decrease from 1997 to 1999, there was an increasing trend in public coverage from 1999 to early 2005. There was an increase from 1997 to 1999 followed by a decreasing trend from 1999 to early 2005 in private coverage for this age group. These trends were observed regardless of whether Method 1 or Method 2 was used to calculate the 2004 estimate, but the extent of the increase or decrease is larger using one method compared to the other method.

**Figure 1.2. Percentage of persons under age 65 years without health insurance coverage, by age group and sex: United States, January–March 2005**



NOTES: A person was defined as uninsured if he or she did not have any private health insurance, Medicare, Medicaid, State Children's Health Insurance Program (SCHIP), State-sponsored or other government-sponsored health plan, or military plan at the time of the interview. A person was also defined as uninsured if he or she had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care. The analyses excluded 189 persons (1.0%) with unknown health insurance status. The data on health insurance status were edited using an automated system based on logic checks and keyword searches. The resulting estimates of persons not having health insurance coverage are generally 0.1–0.2 percentage points lower than those based on the editing procedures used for the final data files.


In quarter 3 of 2004, two additional questions were added to the NHIS insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1 in the data table) and with the responses to these new questions (presented in the figure above and noted as Method 2 in the data table). Estimates for this figure in previous Early Releases were calculated using Method 1 and are, therefore, not directly comparable to estimates in the above figure. Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For both sexes combined, the percentage of those who were uninsured was highest among persons aged 18–24 years (29.7%) and lowest among persons under age 18 years (8.6%) and at ages 45–64 years (13.0%). Starting at age 18 years, younger adults were more likely than older adults to lack health insurance coverage.

■ Among adults in the age groups 18–24 years, 25–34 years, and 35–44 years, men were more likely than women to lack health insurance coverage.

**Figure 1.3. Age-sex-adjusted percentage of persons of all ages without health insurance coverage, by race/ethnicity: United States, January–March 2005**



NOTES: A person was defined as uninsured if he or she did not have any private health insurance, Medicare, Medicaid, State Children's Health Insurance Program (SCHIP), State-sponsored or other government-sponsored health plan, or military plan at the time of the interview. A person was also defined as uninsured if he or she had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care. The analyses excluded 207 persons (1.0%) with unknown health insurance status. The data on health insurance status were edited using an automated system based on logic checks and keyword searches. The resulting estimates of persons not having health insurance coverage are generally 0.1–0.2 percentage points lower than those based on the editing procedures used for the final data files. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

In quarter 3 of 2004, two additional questions were added to the NHIS insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1 in the data table) and with the responses to these new questions (presented in the figure above and noted as Method 2 in the data table). Estimates for this figure in previous Early Releases were calculated using Method 1 and are, therefore, not directly comparable to estimates in the above figure. Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ After adjusting for age and sex, the percentage of uninsured persons was 29.4% for Hispanic persons, 10.2% for non-Hispanic white persons, and 17.9% for non-Hispanic black persons.

■ Hispanic persons were most likely to be uninsured, followed by non-Hispanic black persons and non-Hispanic white persons.

## Data tables for figures 1.1–1.3:

**Data table for figure 1.1. Percentage of persons of all ages without health insurance coverage: United States, 1997–2005**

Year	Percent	95% confidence interval
1997	15.4	15.0-15.8
1998	14.6	14.1-15.1
1999	14.2	13.8-14.6
2000	14.9	14.5-15.3
2001	14.3	13.8-14.8
2002	14.7	14.3-15.1
2003	15.2	14.8-15.7
2004 (Method 1 <sup>1</sup> )	14.7	14.3-15.2
2004 (Method 2 <sup>1</sup> )	14.6	14.2-15.0
January–March 2005	14.4	13.5-15.3

<sup>1</sup>In quarter 3 of 2004, two additional questions were added to the NHIS Insurance section to reduce potential errors in reporting Medicare and Medicaid status. Persons aged 65 years and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Depending on responses to these two questions, respondents may have been reclassified. Estimates of uninsurance for 2004 are calculated both without using the additional information from these new questions (noted as Method 1) and with the responses to these new questions (noted as Method 2). Beginning in 2005, all estimates are calculated using Method 2. See "About This Release" for additional information.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



**Data table for figure 1.2. Percentage of persons under age 65 years without health insurance coverage, by age group and sex: United States, January–March 2005**

<b>Age and sex</b>	<b>Percent</b>	<b>95% confidence interval</b>
<b>Under 18 years</b>		
Total	8.6	7.5-9.8
Male	8.9	7.6-10.2
Female	8.2	6.9-9.8
<b>18–24 years</b>		
Total	29.7	26.8-32.7
Male	35.4	31.7-39.3
Female	23.8	20.4-27.7
<b>25–34 years</b>		
Total	25.5	23.3-27.8
Male	28.9	26.0-31.9
Female	22.2	19.8-24.8
<b>35–44 years</b>		
Total	17.4	15.8-19.1
Male	21.0	18.8-23.4
Female	13.9	12.1-15.8
<b>45–64 years</b>		
Total	13.0	11.9-14.2
Male	13.5	12.1-15.1
Female	12.5	11.2-13.9
<b>Under 65 years: crude<sup>1</sup></b>		
Total	16.2	15.2-17.3
Male	18.2	17.1-19.4
Female	14.3	13.2-15.4
<b>Under 65 years: age-adjusted<sup>2</sup></b>		
Total	16.4	15.4-17.4
Male	18.5	17.3-19.6
Female	14.3	13.3-15.5

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are for persons under 65 years of age and are age-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–44 years, and 45–64 years.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 1.3. Adjusted percentage of persons of all ages without health insurance coverage, by race/ethnicity: United States, January–March 2005**


<b>Race/ethnicity</b>	<b>Age-sex-adjusted<sup>1</sup></b>	<b>Age-adjusted<sup>2</sup></b>
<b>Hispanic or Latino</b>	29.4 (27.1-31.7)	29.0 (26.8-31.3)
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	10.2 (9.2-11.2)	10.5 (9.5-11.7)
<b>Black, single race</b>	17.9 (15.6-20.5)	17.8 (15.5-20.4)

<sup>1</sup>Age-sex-adjusted estimates are presented in the figure. Estimates are for persons of all ages and are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are for persons under 65 years of age and are age-adjusted to the 2000 projected U.S. standard population using four age groups: under 18 years, 18–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 2.1. Percentage of persons of all ages with a usual place to go for medical care: United States, 1997–2005**




NOTES: The usual place to go for medical care does not include a hospital emergency room. The analyses excluded persons with an unknown usual place to go for medical care (about 0.6% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Combined Sample Adult and Sample Child Core components of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, the percentage of persons who had a usual place to go for medical care was 86.7% (95% confidence interval = 85.7%–87.7%), which was similar to the 2004 estimate of 86.8%.
- The annual percentage of persons of all ages who had a usual place to go for medical care was 86.2% in 1997, 87.4% in 1998, 86.3% in 1999, 87.0% in 2000, 88.0% in 2001, 87.6% in 2002, 87.9% in 2003, and 86.8% in 2004.

**Figure 2.2. Percentage of persons of all ages with a usual place to go for medical care, by age group and sex: United States, January–March 2005**




NOTES: The usual place to go for medical care does not include a hospital emergency room. The analyses excluded 66 persons (0.7%) with an unknown usual place to go for medical care.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- Of the five age groups, persons aged 18–24 years were least likely to have a usual place to go for medical care. Children under age 18 years were more likely than adults aged 18–64 years to have a usual place to go for medical care.
- For adults aged 18–64 years, men were less likely than women to have a usual place to go for medical care.
- Among those aged 18 years and over, the percentage of persons having a usual place to go for medical care increased with age.

**Figure 2.3. Age-sex-adjusted percentage of persons of all ages with a usual place to go for medical care, by race/ethnicity: United States, January–March 2005**



NOTES: The usual place to go for medical care does not include a hospital emergency room. The analyses excluded 66 persons (0.7%) with an unknown usual place to go for medical care. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- After adjusting for age and sex, the percentage of persons with a usual place to go for medical care was 78.4% for Hispanic persons, 88.9% for non-Hispanic white persons, and 84.3% for non-Hispanic black persons.
- Of the three race/ethnicity groups, Hispanic persons were least likely to have a usual place to go for medical care.

## Data tables for figures 2.1–2.3:

Data table for figure 2.1. Percentage of persons of all ages with a usual place to go for medical care: United States, 1997–2005

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	86.2 (85.8-86.7)	86.3 (85.9-86.7)
1998	87.4 (86.9-87.8)	87.4 (87.0-87.9)
1999	86.3 (85.9-86.7)	86.4 (86.0-86.8)
2000	87.0 (86.5-87.4)	87.0 (86.6-87.4)
2001	88.0 (87.6-88.5)	88.0 (87.6-88.5)
2002	87.6 (87.1-88.0)	87.6 (87.1-88.0)
2003	87.9 (87.4-88.4)	87.9 (87.4-88.4)
2004	86.8 (86.3-87.3)	86.8 (86.3-87.2)
January–March 2005	86.7 (85.7-87.7)	86.6 (85.6-87.6)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 2.2. Percentage of persons of all ages with a usual place to go for medical care, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>Under 18 years</b>		
Total	95.1	94.1-96.0
Male	94.9	93.6-96.0
Female	95.3	93.6-96.7
<b>18–24 years</b>		
Total	68.9	63.8-73.6
Male	58.4	50.1-66.1
Female	79.6	74.4-83.9
<b>25–44 years</b>		
Total	78.1	76.1-80.0
Male	69.7	66.5-72.7
Female	86.4	84.1-88.4
<b>45–64 years</b>		
Total	90.3	88.9-91.5
Male	88.1	86.0-90.0
Female	92.3	90.5-93.8
<b>65 years and over</b>		
Total	96.5	95.3-97.5
Male	97.2	95.3-98.4
Female	96.0	94.1-97.3
<b>All ages: crude<sup>1</sup></b>		
Total	86.7	85.7-87.7
Male	82.6	81.0-84.2
Female	90.7	89.5-91.7
<b>All ages: age-adjusted<sup>2</sup></b>		
Total	86.6	85.6-87.6
Male	82.7	81.1-84.1
Female	90.6	89.4-91.6

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 2.3. Adjusted percentage of persons of all ages with a usual place to go for medical care, by race/ethnicity: United States, January–March 2005**

Race/ethnicity	Percent (95% confidence interval)	
	Age-sex-adjusted <sup>1</sup>	Age-adjusted <sup>2</sup>
<b>Hispanic or Latino</b>	78.4 (76.2-80.4)	78.1 (75.8-80.1)
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	88.9 (87.7-90.0)	88.9 (87.7-90.1)
<b>Black, single race</b>	84.3 (81.5-86.8)	84.7 (81.9-87.1)


<sup>1</sup>Age-sex-adjusted estimates are presented in the figure. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



**Figure 3.1. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months: United States, 1997–2005**




NOTES: The analyses excluded persons with unknown success in obtaining needed medical care (about 0.5% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 5.7% (95% confidence interval = 5.2%–6.2%) of the population was unable to obtain needed medical care due to cost at some time during the past 12 months. The estimate is slightly higher than, but not significantly different from, the 2004 estimate of 5.5%.

■ From 1998 through early 2005, an increasing trend has been observed in the percentage of persons who experienced this lack of access to medical care (4.2% in 1998 to 5.7% in early 2005).

**Figure 3.2. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by age group and sex: United States, January–March 2005**




NOTE: The analyses excluded 53 persons (0.3%) with unknown success in obtaining needed medical care.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who were unable to obtain needed medical care due to cost at some time during the past 12 months was 2.4% for those aged under 18 years, 7.6% for those aged 18–64 years, and 2.9% for those aged 65 years and over.
- For adults 18–64 years, women were more likely than men to have experienced this lack of access to medical care during the past 12 months.

**Figure 3.3. Age-sex-adjusted percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by race/ethnicity: United States, January–March 2005**



NOTES: The analyses excluded 53 persons (0.3%) with unknown success in obtaining needed medical care. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- After adjusting for age and sex, 5.7% of Hispanic persons, 5.5% of non-Hispanic white persons, and 7.2% of non-Hispanic black persons were unable to obtain needed medical care due to cost at some time during the year preceding the interview.
- Non-Hispanic black persons were more likely than non-Hispanic white persons to experience this lack of access to medical care.

## Data tables for figures 3.1–3.3:

**Data table for figure 3.1. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months: United States, 1997–2005**

Year	Percent	95% confidence interval
1997	4.5	4.3-4.7
1998	4.2	4.0-4.4
1999	4.3	4.1-4.5
2000	4.5	4.3-4.7
2001	4.7	4.5-4.9
2002	4.7	4.5-4.9
2003	5.3	5.1-5.5
2004	5.5	5.2-5.7
January–March 2005	5.7	5.2-6.2

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 3.2. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>Under 18 years</b>		
Total	2.4	1.9-3.1
Male	2.9	2.1-3.9
Female	2.0	1.4-2.9
<b>18–64 years</b>		
Total	7.6	6.9-8.2
Male	6.5	5.8-7.3
Female	8.6	7.8-9.5
<b>65 years and over</b>		
Total	2.9	2.2-3.9
Male	3.1	2.1-4.5
Female	2.8	2.0-3.8
<b>All ages: crude<sup>1</sup></b>		
Total	5.7	5.2-6.2
Male	5.2	4.6-5.8
Female	6.2	5.7-6.8
<b>All ages: age-adjusted<sup>2</sup></b>		
Total	5.7	5.2-6.2
Male	5.1	4.6-5.7
Female	6.2	5.6-6.8

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 3.3. Age-sex-adjusted percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by race/ethnicity: United States, January–March 2005**

<b>Race/ethnicity</b>	<b>Percent<sup>1</sup></b>	<b>95% confidence interval</b>
<b>Hispanic or Latino</b>	5.7	4.7-6.7
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	5.5	4.9-6.1
<b>Black, single race</b>	7.2	5.8-8.8

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 4.1. Percentage of adults aged 18 years and over who had received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–2005**



NOTES: From 1997 to August 2003, respondents were asked if they had received a flu shot during the past 12 months. Beginning in September 2003, respondents were asked about influenza vaccination by nasal spray (sometimes called by the brand name FluMist™) during the past 12 months in addition to the question regarding the flu shot. Beginning in 2005, receipt of nasal spray flu vaccinations was included in the calculation of flu vaccination estimates. The impact of this change on the estimates was minimal (typically 0.5 or less). Responses to the flu vaccination questions cannot be used to determine when during the preceding 12 months the subject received the flu vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of a flu vaccination is seasonal. According to the recommendations of the Advisory Committee on Immunization Practices, all adults aged 65 years and over should receive an influenza vaccination (6). In the 2000–01 flu season (but not implemented until the 2001–02 flu season), these recommendations for influenza vaccination were expanded to include persons 50 to 64 years of age, a group for which influenza vaccination was formerly recommended only if they had existing high-risk conditions (7). Adults aged 18–49 years are recommended to receive influenza vaccination only if they have existing high-risk conditions. The analyses excluded those with unknown flu vaccination status (about 1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In the first quarter of 2005, the percentage of adults who had received an influenza vaccination during the past year was 59.6% for persons aged 65 years and over, 21.2% for persons aged 50–64 years, and 10.2% for persons aged 18–49 years.

■ For all three age groups, first quarter estimates in 2005 were considerably lower than first quarter estimates in 2004. An influenza vaccination shortage occurred during the 2004–05 flu season (6). Previous delays in availability of the flu shots occurred in the fall of 2000 and, to a lesser extent, in the fall of 2001 (7,8).

**Table 4.1. Annual percentage of adults aged 50 years and over who had received an influenza shot during the past 12 months, by age group and sex: United States, 1997–2004**

Year	50–64 years			65 years and over			
	Total	Men	Women	Total		Men	Women
				Crude	Age-adjusted		
1997	31.9 (30.5-33.3)	28.0 (26.1-29.9)	35.5 (33.6-37.4)	63.2 (61.9-64.6)	63.1 (61.7-64.4)	64.8 (62.5-67.1)	62.1 (60.5-63.7)
1998	33.1 (31.7-34.5)	29.0 (27.0-31.0)	37.0 (35.1-38.9)	63.3 (61.9-64.7)	63.3 (61.9-64.6)	63.7 (61.5-65.9)	63.0 (61.2-64.8)
1999	34.1 (32.8-35.4)	30.5 (28.6-32.4)	37.4 (35.5-39.3)	65.7 (64.3-67.2)	65.1 (63.6-66.5)	67.2 (65.0-69.4)	64.6 (62.7-66.5)
2000	34.6 (33.1-36.1)	31.9 (29.9-33.9)	37.2 (35.2-39.1)	64.4 (63.0-65.9)	64.6 (63.2-66.0)	66.0 (63.8-68.3)	63.3 (61.6-65.0)
2001	32.2 (30.9-33.5)	30.3 (28.3-32.2)	34.0 (32.2-35.8)	63.1 (61.7-64.5)	63.2 (61.8-64.6)	64.8 (62.5-67.1)	61.8 (60.1-63.5)
2002	34.0 (32.7-35.3)	30.7 (28.8-32.5)	37.2 (35.4-38.9)	65.7 (64.3-67.2)	65.9 (64.5-67.3)	67.1 (64.7-69.5)	64.7 (62.8-66.6)
2003	36.8 (35.4-38.2)	34.5 (32.6-36.3)	38.9 (37.0-40.9)	65.5 (64.1-66.9)	65.6 (64.2-66.9)	66.0 (63.9-68.1)	65.1 (63.2-67.0)
2004	35.9 (34.6-37.3)	33.3 (31.3-35.3)	38.5 (36.7-40.3)	64.6 (63.2-66.1)	64.7 (63.2-66.1)	64.1 (61.9-66.3)	65.0 (63.3-66.7)


NOTES: From 1997 to August 2003, respondents were asked if they had received a flu shot during the past 12 months. Beginning in September 2003, respondents were asked about influenza vaccination by nasal spray (sometimes called by the brand name FluMist™) during the past 12 months in addition to the question regarding the flu shot. Receipt of nasal spray flu vaccinations was not included in the calculation of flu vaccination estimates for Early Releases before 2005. Responses to the flu vaccination questions cannot be used to determine when during the preceding 12 months the subject received the flu vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of a flu shot is seasonal. According to the recommendations of the Advisory Committee on Immunization Practices, all adults aged 65 years and over should receive an influenza vaccination (6). In the 2000–01 flu season (but not implemented until the 2001–02 flu season), these recommendations for influenza vaccination were expanded to include persons 50 to 64 years of age, a group for which influenza vaccination was formerly recommended only if they had existing high-risk conditions (7). Adults aged 18–49 years are recommended to receive influenza vaccination only if they have existing high-risk conditions. The analyses excluded those with unknown flu shot status (about 1% of respondents each year). Age-adjusted estimates for persons aged 65 years and over for this Healthy People 2010 Leading Health Indicator are based on the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2004 National Health Interview Surveys. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For adults aged 50–64 years, the annual percentage of persons who received a flu shot during the past 12 months was 35.9% in 2004. The estimate in 2004 was lower than, but not significantly different from, that in 2003. This pattern was seen in men and women.

■ For adults aged 65 years and over, the annual percentage of persons who received a flu shot was 64.6% in 2004. The estimate in 2004 was lower than, but not significantly different from, that in 2003. This pattern was seen in men and women.

**Figure 4.2. Percentage of persons who had received an influenza vaccination during the past 12 months, by age group and sex: United States, January–March 2005**




NOTES: Respondents were asked about receipt of influenza vaccination by nasal spray (sometimes called by the brand name FluMist™) during the past 12 months in addition to a question regarding receipt of a flu shot during the past 12 months. Responses to these questions cannot be used to determine when during the preceding 12 months the subject received the flu vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of a flu vaccination is seasonal. According to the recommendations of the Advisory Committee on Immunization Practices, all adults aged 65 years and over should receive an influenza vaccination (6). In the 2000–01 flu season (but not implemented until the 2001–02 flu season), these recommendations for influenza vaccination were expanded to include persons 50 to 64 years of age, a group for which influenza vaccination was formerly recommended only if they had existing high-risk conditions (7). The recommendations were again expanded for the 2003–2004 flu season to include children aged 6–23 months (9). Persons aged 2–49 years are recommended to receive an influenza vaccination only if they have existing high-risk conditions. The analyses excluded 133 persons (1.4%) with unknown flu vaccination status. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who had a flu vaccination during the past 12 months was highest among persons aged 65 years and over (59.6%), followed by persons aged 50–64 years (21.2%), 0–17 years (16.1%), and 18–49 years (10.2%).
- For adults aged 18–49 years, women were more likely than men to have received a flu vaccination during the past 12 months.



**Figure 4.3. Percentage of adults aged 65 years and over who had received an influenza vaccination during the past 12 months, by race/ethnicity: United States, January–March 2005**



NOTES: Respondents were asked about receipt of influenza vaccination by nasal spray (sometimes called by the brand name FluMist™) during the past 12 months in addition to a question regarding receipt of a flu shot during the past 12 months. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of a flu vaccination is seasonal. The analyses excluded 22 adults (1.7%) aged 65 years and over with unknown flu vaccination status. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For adults aged 65 years and over, the percentage of persons receiving a flu vaccination during the past 12 months was 44.9% for Hispanic persons, 63.3% for non-Hispanic white persons, and 39.0% for non-Hispanic black persons.
- Hispanic persons and non-Hispanic black persons were less likely than non-Hispanic white persons to have received a flu vaccination during the past 12 months.

## Data tables for figures 4.1–4.3:

Data table for figure 4.1. Percentage of adults aged 18 years and over who had received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–2005

Year and quarter	Percent (95% confidence interval)		
	18–49 years	50–64 years	65 years and over
<b>1997</b>			
Quarter 1	13.6 (12.5-14.6)	30.5 (27.8-33.2)	65.0 (62.3-67.6)
Quarter 2	14.5 (13.4-15.5)	31.3 (28.7-34.0)	63.7 (61.1-66.2)
Quarter 3	13.6 (12.6-14.6)	32.0 (29.3-34.6)	63.1 (60.3-65.9)
Quarter 4	15.6 (14.5-16.7)	33.6 (31.1-36.2)	61.2 (58.7-63.8)
<b>1998</b>			
Quarter 1	16.1 (14.8-17.3)	34.2 (31.3-37.1)	66.3 (63.2-69.4)
Quarter 2	15.3 (14.1-16.5)	32.8 (30.1-35.5)	64.0 (61.3-66.8)
Quarter 3	14.5 (13.3-15.6)	32.0 (29.3-34.6)	61.3 (58.5-64.0)
Quarter 4	16.0 (14.8-17.2)	33.5 (30.8-36.1)	61.6 (58.7-64.5)
<b>1999</b>			
Quarter 1	16.3 (14.8-17.7)	34.2 (31.1-37.3)	67.0 (64.0-70.1)
Quarter 2	16.0 (14.7-17.3)	32.6 (29.8-35.4)	66.1 (63.4-68.8)
Quarter 3	15.8 (14.5-17.1)	32.8 (30.1-35.5)	64.1 (61.2-67.0)
Quarter 4	17.6 (16.2-18.9)	36.7 (34.2-39.2)	65.7 (62.7-68.6)
<b>2000</b>			
Quarter 1	18.6 (17.2-19.9)	36.6 (33.7-39.4)	68.2 (65.3-71.0)
Quarter 2	18.0 (16.7-19.4)	38.5 (35.7-41.4)	68.1 (65.6-70.7)
Quarter 3	18.2 (16.9-19.4)	36.6 (33.7-39.5)	67.1 (64.4-69.8)
Quarter 4	13.6 (12.4-14.8)	26.6 (24.2-29.0)	54.3 (51.6-57.1)
<b>2001</b>			
Quarter 1	14.7 (13.4-16.0)	32.3 (29.6-35.0)	63.3 (60.2-66.3)
Quarter 2	15.9 (14.7-17.1)	32.6 (30.1-35.1)	65.4 (62.8-68.0)
Quarter 3	14.9 (13.9-15.9)	33.3 (30.7-35.8)	64.0 (61.1-66.8)
Quarter 4	14.5 (13.6-15.9)	30.6 (28.0-33.1)	59.6 (56.7-62.4)
<b>2002</b>			
Quarter 1	16.4 (15.2-17.7)	36.3 (33.6-38.9)	66.6 (63.8-69.4)
Quarter 2	16.0 (14.8-17.2)	33.0 (30.5-35.5)	67.8 (65.3-70.3)
Quarter 3	16.2 (14.9-17.5)	33.1 (30.6-35.6)	63.1 (60.5-65.8)
Quarter 4	16.4 (15.1-17.8)	33.8 (31.0-36.6)	65.5 (62.4-68.6)
<b>2003</b>			
Quarter 1	17.1 (15.7-18.4)	36.8 (34.2-39.4)	67.8 (65.0-70.6)
Quarter 2	17.2 (15.8-18.6)	36.8 (33.9-39.7)	65.4 (62.6-68.3)
Quarter 3	16.4 (15.2-17.6)	37.4 (34.9-39.9)	65.4 (62.8-67.9)
Quarter 4	16.7 (15.2-18.1)	36.1 (33.3-39.0)	63.3 (60.1-66.5)
<b>2004</b>			
Quarter 1	19.3 (17.9-20.8)	40.6 (38.0-43.3)	70.3 (67.5-73.0)
Quarter 2	20.9 (19.1-22.6)	40.0 (37.1-43.0)	69.5 (66.7-72.3)
Quarter 3	19.4 (18.2-20.7)	41.0 (38.4-43.6)	66.4 (63.6-69.2)
Quarter 4	12.0 (10.9-13.1)	22.3 (20.2-24.5)	52.4 (49.5-55.4)
<b>2005</b>			
Quarter 1	10.2 (9.1-11.5)	21.2 (19.1-23.4)	59.6 (56.5-62.7)

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 4.2. Percentage of persons who had received an influenza vaccination during the past 12 months, by age group and sex: United States, January–March 2005**

<b>Age and sex</b>	<b>Percent</b>	<b>95% confidence interval</b>
<b>0–2 years</b>		
Total	35.9	31.0-41.1
Male	37.1	30.2-44.7
Female	34.5	27.6-42.2
<b>3–4 years</b>		
Total	22.5	17.2-28.8
Male	21.7	14.8-30.7
Female	23.2	16.2-32.0
<b>5–11 years</b>		
Total	12.9	10.5-15.6
Male	13.5	10.3-17.5
Female	12.3	9.2-16.2
<b>12–17 years</b>		
Total	8.1	6.2-10.5
Male	7.3	5.1-10.3
Female	9.0	6.0-13.3
<b>18–49 years</b>		
Total	10.2	9.1-11.5
Male	7.7	6.4-9.3
Female	12.6	11.0-14.5
<b>50–64 years</b>		
Total	21.2	19.1-23.4
Male	20.0	16.9-23.5
Female	22.3	19.5-25.4
<b>65 years and over</b>		
Total	59.6	56.5-62.7
Male	58.3	53.5-63.0
Female	60.6	56.4-64.6
<b>0–17 years</b>		
Total	16.1	14.4-18.0
Male	16.2	14.0-18.8
Female	16.0	13.7-18.7
<b>18 years and over: crude<sup>1</sup></b>		
Total	20.6	19.4-21.8
Male	17.7	16.2-19.3
Female	23.3	21.6-25.1
<b>65 years and over: age-adjusted<sup>2</sup></b>		
Total	59.7	56.6-62.7
Male	59.3	54.8-63.6
Female	60.3	56.1-64.4

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 4.3. Percentage of adults aged 65 years and over who had received an influenza vaccination during the past 12 months, by race/ethnicity: United States, January–March 2005**


Race/ethnicity	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
<b>Hispanic or Latino</b>	44.9 (32.8-57.6)	45.2 (33.3-57.7)
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	63.3 (59.9-66.5)	63.1 (59.8-66.3)
<b>Black, single race</b>	39.0 (30.5-48.1)	38.8 (30.2-48.1)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 5.1. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination: United States, 1997–2005**




NOTES: The analyses excluded those with unknown pneumococcal vaccination status (about 3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, the percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination was 56.6% (95% confidence interval = 53.3%–59.7%), which was similar to the 2004 estimate of 56.8%.

■ For adults aged 65 years and over, there was an increasing trend in the percentage of those who had ever received a pneumococcal vaccination from 1997 to 2004.

**Figure 5.2. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–March 2005**




NOTES: The analyses excluded 49 adults (3.7%) aged 65 years and over with unknown pneumococcal vaccination status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The percentage of adults who had ever received a pneumococcal vaccination was 50.5% for persons aged 65–74 years and 63.5% for persons aged 75 years and over.

■ For adults aged 65–74 years, women were more likely than men to have ever received a pneumococcal vaccination.

**Figure 5.3. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–March 2005**



NOTE: The analyses excluded 49 adults (3.7%) aged 65 years and over with unknown pneumococcal vaccination status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination was 22.2% for Hispanic persons, 61.6% for non-Hispanic white persons, and 36.7% for non-Hispanic black persons.

- Hispanic persons and non-Hispanic black persons were less likely than non-Hispanic white persons to have ever received a pneumococcal vaccination.

## Data tables for figures 5.1–5.3:

**Data table for figure 5.1. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination: United States, 1997–2005**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	42.4 (40.9-43.9)	42.6 (41.1-44.1)
1998	46.0 (44.5-47.5)	46.3 (44.8-47.8)
1999	49.7 (48.1-51.3)	49.9 (48.3-51.5)
2000	53.1 (51.5-54.7)	53.4 (51.8-55.0)
2001	54.0 (52.4-55.6)	54.2 (52.6-55.7)
2002	56.0 (54.5-57.5)	56.2 (54.8-57.7)
2003	55.6 (54.1-57.1)	55.7 (54.2-57.1)
2004	56.8 (55.3-58.3)	57.0 (55.5-58.5)
January–March 2005	56.6 (53.3-59.7)	56.7 (53.5-59.8)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 5.2. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>65–74 years</b>		
Total	50.5	46.2-54.8
Male	44.0	38.0-50.2
Female	56.2	50.5-61.8
<b>75 years and over</b>		
Total	63.5	59.3-67.4
Male	65.3	57.9-72.0
Female	62.3	57.0-67.3
<b>65 years and over: crude<sup>1</sup></b>		
Total	56.6	53.3-59.7
Male	53.1	48.1-57.9
Female	59.2	55.2-63.1
<b>65 years and over: age-adjusted<sup>2</sup></b>		
Total	56.7	53.5-59.8
Male	54.2	49.4-58.9
Female	59.1	55.1-63.0

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



**Data table for figure 5.3. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–March 2005**


Race/ethnicity	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
<b>Hispanic or Latino</b>	22.2 (13.9-33.5)	23.2 (15.0-34.2)
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	61.6 (58.1-64.9)	61.5 (58.0-64.8)
<b>Black, single race</b>	36.7 (29.1-45.0)	36.3 (28.7-44.7)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–2005**




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m<sup>2</sup> or more. The measure is based on self-reported height and weight. The analyses excluded people with unknown height or weight (about 4% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, 25.1% (95% confidence interval = 23.8%–26.5%) of U.S. adults aged 20 years and over were obese, which was similar to the 2004 estimate of 24.5%.
- The prevalence of obesity among U.S. adults aged 20 years and over has increased over time from 19.4% in 1997 to 25.1% in early 2005. The estimate in 2003 was lower than, but not significantly different from, that in 2002.

**Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–March 2005**




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m<sup>2</sup> or more. The measure is based on self-reported height and weight. The analyses excluded 309 people (4.5%) with unknown height or weight.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of obesity was highest among adults aged 40–59 years (29.0%) and lowest among adults aged 20–39 years (21.7%). This pattern in obesity by age group was seen in both men and women.
- There was no significant difference in the prevalence of obesity between women and men for all three age groups.

**Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–March 2005**



NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m<sup>2</sup> or more. The measure is based on self-reported height and weight. The analyses excluded 309 people (4.5%) with unknown height or weight. Estimates are age-adjusted to the 2000 projected U.S. standard population using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ Non-Hispanic white women were less likely than Hispanic and non-Hispanic black women to be obese. This race/ethnicity difference was not seen among men.

■ Among the six sex-specific race/ethnicity groups, the age-adjusted prevalence of obesity was highest among non-Hispanic black women (36.5%).

## Data tables for figures 6.1–6.3:

**Data table for figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–2005**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	19.4 (18.9-19.9)	19.5 (18.9-20.0)
1998	20.6 (20.1-21.1)	20.6 (20.0-21.1)
1999	21.5 (20.9-22.1)	21.5 (20.9-22.1)
2000	21.8 (21.2-22.4)	21.8 (21.2-22.3)
2001	23.0 (22.4-23.6)	22.9 (22.3-23.5)
2002	23.9 (23.3-24.6)	23.8 (23.2-24.5)
2003	23.7 (23.1-24.3)	23.5 (22.9-24.2)
2004	24.5 (23.9-25.1)	24.3 (23.8-25.0)
January–March 2005	25.1 (23.8-26.5)	25.0 (23.7-26.3)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>20–39 years</b>		
Total	21.7	19.8-23.8
Male	21.4	18.6-24.4
Female	22.0	19.5-24.7
<b>40–59 years</b>		
Total	29.0	26.8-31.3
Male	29.5	26.5-32.6
Female	28.5	25.6-31.7
<b>60 years and over</b>		
Total	24.2	21.9-26.5
Male	25.0	21.2-29.2
Female	23.5	20.6-26.7
<b>20 years and over: crude<sup>1</sup></b>		
Total	25.1	23.8-26.5
Male	25.4	23.4-27.4
Female	24.9	23.1-26.7
<b>20 years and over: age-adjusted<sup>2</sup></b>		
Total	25.0	23.7-26.3
Male	25.1	23.2-27.2
Female	24.8	23.0-26.6

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–March 2005**

Sex and race/ethnicity	Percent <sup>1</sup>	95% confidence interval
<b>Male</b>		
Hispanic or Latino	24.1	19.7-29.0
Not Hispanic or Latino:		
White, single race	24.7	22.4-27.1
Black, single race	29.7	24.6-35.4
<b>Female</b>		
Hispanic or Latino	31.0	26.2-36.4
Not Hispanic or Latino:		
White, single race	21.3	19.3-23.4
Black, single race	36.5	31.7-41.7

<sup>1</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–2005**




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator (Healthy People 2010). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. The analyses excluded persons with unknown physical activity participation (about 3% of respondents each year). In Early Releases before June 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than that of Healthy People 2010. The earlier ER estimates excluded from the analysis persons with unknown duration who were known to have not met the frequency recommendations (i.e. partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. All estimates have been rerun using the revised denominator. The impact of the change on the estimates was minimal (typically 0.1 or less). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- In early 2005, 30.8% (95% confidence interval = 29.2%–32.4%) of U.S. adults aged 18 years and over engaged in regular leisure-time physical activity. This estimate was similar to the 2004 quarter one estimate of 29.6%.
- The annual percentages of adults aged 18 years and over who engaged in regular leisure-time physical activity increased from 29.8% in 1998 to 31.9% in 2000, did not change significantly from 2000 to 2003, and decreased from 2003 to 2004.

**Figure 7.2. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January–March 2005**




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator (Healthy People 2010). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before June 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than that of Healthy People 2010. The earlier ER estimates excluded from the analysis persons with unknown duration who were known to have not met the frequency recommendations (i.e. partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 168 persons (2.4%) with unknown physical activity participation.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of adults who engaged in regular leisure-time physical activity decreased with age.
- For adults 18–24 years, women were less likely than men to engage in regular leisure-time physical activity.



**Figure 7.3. Age-sex-adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January–March 2005**



NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator (Healthy People 2010). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before June 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than that of Healthy People 2010. The earlier ER estimates excluded from the analysis persons with unknown duration who were known to have not met the frequency recommendations (i.e. partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 168 persons (2.4%) with unknown physical activity participation. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted percentage of adults who engaged in regular leisure-time physical activity was 21.3% for Hispanic adults, 34.8% for non-Hispanic white adults, and 19.2% for non-Hispanic black adults.

■ Non-Hispanic white adults were more likely to engage in regular leisure-time physical activity than Hispanic adults and non-Hispanic black adults.

## Data tables for figures 7.1–7.3:

Data table for figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–2005

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
<b>1997 Yearly</b>	32.0 (31.3-32.6)	31.8 (31.1-32.5)
<b>January–March</b>	31.3 (30.1-32.6)	31.1 (29.9-32.5)
<b>1998 Yearly</b>	29.8 (29.1-30.4)	29.6 (28.9-30.3)
<b>January–March</b>	28.8 (27.4-30.1)	28.5 (27.2-29.9)
<b>1999 Yearly</b>	30.2 (29.5-31.0)	30.1 (29.4-30.8)
<b>January–March</b>	27.6 (26.2-29.1)	27.4 (26.0-28.9)
<b>2000 Yearly</b>	31.9 (31.2-32.6)	31.8 (31.1-32.5)
<b>January–March</b>	31.5 (30.1-32.9)	31.3 (30.0-32.7)
<b>2001 Yearly</b>	32.1 (31.3-32.8)	31.9 (31.2-32.7)
<b>January–March</b>	30.6 (29.2-32.0)	30.5 (29.1-31.9)
<b>2002 Yearly</b>	31.9 (31.1-32.7)	31.8 (31.0-32.6)
<b>January–March</b>	31.5 (30.0-33.0)	31.4 (30.0-32.9)
<b>2003 Yearly</b>	32.8 (32.1-33.6)	32.8 (32.0-33.6)
<b>January–March</b>	31.2 (29.7-32.6)	31.1 (29.7-32.5)
<b>2004 Yearly</b>	30.2 (29.4-31.0)	30.1 (29.4-30.9)
<b>January–March</b>	29.6 (28.2-31.0)	29.5 (28.1-30.9)
<b>2005 January–March</b>	30.8 (29.2-32.4)	30.7 (29.2-32.3)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 7.2. Percentage of adults aged 18 and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January–March 2005**

<b>Age and sex</b>	<b>Percent</b>	<b>95% confidence interval</b>
<b>18–24 years</b>		
Total	39.7	35.0-44.7
Male	45.0	37.0-53.1
Female	34.3	29.1-40.0
<b>25–64 years</b>		
Total	31.2	29.4-33.0
Male	31.2	28.7-33.7
Female	31.2	28.9-33.6
<b>65–74 years</b>		
Total	25.0	21.4-29.1
Male	27.5	22.3-33.3
Female	23.0	18.1-28.8
<b>75 years and over</b>		
Total	18.0	14.5-22.1
Male	21.3	16.1-27.6
Female	15.9	11.8-21.0
<b>18 years and over: crude<sup>1</sup></b>		
Total	30.8	29.2-32.4
Male	32.2	29.9-34.5
Female	29.5	27.5-31.5
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	30.7	29.2-32.3
Male	31.9	29.8-34.2
Female	29.7	27.7-31.7

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 7.3. Adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January–March 2005**


Race/ethnicity	Percent (95% confidence interval)	
	Age-sex-adjusted <sup>1</sup>	Age-adjusted <sup>2</sup>
<b>Hispanic or Latino</b>	21.3 (18.5-24.3)	21.2 (18.4-24.3)
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	34.8 (33.0-36.7)	34.8 (33.0-36.7)
<b>Black, single race</b>	19.2 (16.3-22.4)	18.9 (16.1-22.2)

<sup>1</sup>Age-sex-adjusted estimates are presented in the figure. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 8.1. Prevalence of current smoking among adults aged 18 years and over: United States, 1997–2005**




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded persons with unknown smoking status (about 1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 22.0% (95% confidence interval = 20.8%–23.4%) of adults aged 18 years and over were current smokers, which was higher than, but not significantly different from, the 2004 estimate of 20.9%.

■ The prevalence of current smoking among U.S. adults declined from 24.7% in 1997 to 20.9% in 2004.

**Figure 8.2. Percent distribution of smoking status among adults aged 18 years and over, by sex: United States, January–March 2005**




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 54 persons (0.8%) with unknown smoking status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The percentage of current smokers was higher for men (26.0%) than for women (18.4%).

■ The percentage of former smokers was higher for men than for women, and the percentage of those who had never smoked was higher for women than for men.

**Figure 8.3. Prevalence of current smoking among adults aged 18 years and over, by age group and sex: United States, January–March 2005**




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 54 persons (0.8%) with unknown smoking status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of adults who were current smokers was lower among adults aged 65 years and over (10.3%) than among adults aged 18–44 years (25.1%) and 45–64 years (23.1%). This pattern in current smoking by age group was seen in both men and women.
- For the age groups 18–44 years and 45–64 years, men were more likely than women to be current smokers.

**Figure 8.4. Age-sex-adjusted prevalence of current smoking among adults aged 18 years and over, by race/ethnicity: United States, January–March 2005**



NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 54 persons (0.8%) with unknown smoking status. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The age-sex-adjusted prevalence of current smoking was 15.0% for Hispanic persons, 23.8% for non-Hispanic white persons, and 24.3% for non-Hispanic black persons.
- Non-Hispanic white adults and non-Hispanic black adults were more likely than Hispanic adults to be current smokers.



## Data tables for figures 8.1–8.4:

**Data table for figure 8.1. Prevalence of current smoking among adults aged 18 years and over: United States, 1997–2005**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	24.7 (24.1-25.3)	24.6 (24.0-25.1)
1998	24.1 (23.5-24.7)	24.0 (23.1-26.8)
1999	23.5 (22.9-24.1)	23.3 (22.7-24.0)
2000	23.2 (22.5-23.8)	23.1 (22.5-23.7)
2001	22.7 (22.1-23.3)	22.6 (22.0-23.2)
2002	22.4 (21.7-23.0)	22.3 (21.7-22.9)
2003	21.6 (21.0-22.2)	21.5 (20.9-22.1)
2004	20.9 (20.3-21.5)	20.8 (20.2-21.4)
January–March 2005	22.0 (20.8-23.4)	22.0 (20.7-23.3)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 8.2. Percent distribution of smoking status among adults aged 18 years and over, by sex: United States, January–March 2005**

Smoking status and sex	Percent	95% confidence interval
<b>Never</b>		
Total	56.9	55.4-58.4
Male	49.6	47.3-51.8
Female	63.7	61.9-65.4
<b>Former</b>		
Total	21.1	20.0-22.2
Male	24.5	22.8-26.2
Female	17.9	16.5-19.5
<b>Current</b>		
Total	22.0	20.8-23.4
Male	26.0	23.9-28.1
Female	18.4	17.0-19.8

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 8.3. Prevalence of current smoking among adults aged 18 years and over, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>18–44 years</b>		
Total	25.1	23.0-27.2
Male	28.4	25.3-31.7
Female	21.8	19.7-24.1
<b>45–64 years</b>		
Total	23.1	21.1-25.1
Male	28.6	25.5-31.9
Female	17.8	15.5-20.4
<b>65 years and over</b>		
Total	10.3	8.6-12.3
Male	10.9	8.3-14.3
Female	9.8	7.7-12.5
<b>18 years and over: crude<sup>1</sup></b>		
Total	22.0	20.8-23.4
Male	26.0	23.9-28.1
Female	18.4	17.0-19.8
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	22.0	20.7-23.3
Male	25.5	23.5-27.6
Female	18.6	17.2-20.0

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 8.4. Adjusted prevalence of current smoking among adults aged 18 years and over, by race/ethnicity: United States, January–March 2005**


Race/ethnicity	Percent (95% confidence interval)	
	Age-sex-adjusted <sup>1</sup>	Age-adjusted <sup>2</sup>
Hispanic or Latino	15.0 (12.8-17.5)	15.2 (13.0-17.8)
<b>Not Hispanic or Latino:</b>		
White, single race	23.8 (22.1-25.5)	23.8 (22.1-25.5)
Black, single race	24.3 (20.9-28.0)	23.9 (20.5-27.6)

<sup>1</sup>Age-sex-adjusted estimates are presented in the figure. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

<sup>2</sup>Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 9.1. Percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year: United States, 1997–2005**




NOTES: The analyses excluded adults with unknown alcohol consumption (about 2% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, the percentage of adults who had five or more drinks in 1 day at least once in the past year was 20.4% (95% confidence interval = 19.0%–21.8%), which was higher than, but not significantly different from, the 2004 estimate of 19.1%.

■ From 2001 through 2003, the percentage of adults who had five or more drinks in 1 day at least once in the past year decreased from 20.0% to 19.1% and remained stable from 2003 to 2004.

**Figure 9.2. Percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year, by age group and sex: United States, January–March 2005**




NOTE: The analyses excluded 133 adults (1.9%) with unknown alcohol consumption.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both men and women, younger adults were more likely than older adults to have had five or more drinks in 1 day at least once in the past year.
- In all four age groups, men were considerably more likely than women to have had five or more drinks in 1 day at least once in the past year.

**Figure 9.3. Age-sex-adjusted percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year, by race/ethnicity: United States, January–March 2005**



NOTES: The analyses excluded 133 adults (1.9%) with unknown alcohol consumption. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using four age groups: 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted percentage of adults who had five or more drinks in 1 day at least once in the past year was 17.3% for Hispanic adults, 23.2% for non-Hispanic white adults, and 12.8% for non-Hispanic black adults.

■ Non-Hispanic white adults were most likely to have had five or more drinks in 1 day at least once in the past year, followed by Hispanic adults and non-Hispanic black adults.

## Data tables for figures 9.1–9.3:

Data table for figure 9.1. Percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year: United States, 1997–2005

Year	Percent	95% confidence interval
1997	21.4	20.8-22.0
1998	20.2	19.6-20.8
1999	20.3	19.6-21.0
2000	19.2	18.6-19.9
2001	20.0	19.4-20.6
2002	19.9	19.2-20.5
2003	19.1	18.5-19.8
2004	19.1	18.4-19.8
January–March 2005	20.4	19.0-21.8

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 9.2. Percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>18–24 years</b>		
Total	35.5	30.7-40.7
Male	46.2	38.6-54.0
Female	24.9	19.7-31.0
<b>25–44 years</b>		
Total	26.5	24.3-28.8
Male	36.7	33.3-40.2
Female	16.7	14.6-19.0
<b>45–64 years</b>		
Total	15.4	13.7-17.4
Male	22.8	19.9-26.0
Female	8.5	6.8-10.6
<b>65 years and over</b>		
Total	3.8	2.7-5.2
Male	6.7	4.7-9.6
Female	-	-
<b>18 years and over: crude<sup>1</sup></b>		
Total	20.4	19.0-21.8
Male	29.0	26.8-31.4
Female	12.4	11.0-13.8
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	20.5	19.2-21.9
Male	28.7	26.6-30.8
Female	12.7	11.4-14.1

-Estimates with a relative standard error greater than 30% are indicated with a dash, but data are not shown.

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using four age groups: 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 9.3. Age-sex-adjusted percentage of adults aged 18 years and over who had 5 or more drinks in 1 day at least once in the past year, by race/ethnicity: United States, January–March 2005**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
Hispanic or Latino	17.3	15.1-19.8
<b>Not Hispanic or Latino:</b>		
White, single race	23.2	21.6-24.9
Black, single race	12.8	10.2-15.9

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using four age groups: 18–24 years, 25–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 10.1. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus: United States, 1997–2005**



NOTES: The analyses excluded those with unknown human immunodeficiency virus (HIV) test status (about 4% of respondents each year). Individuals who received HIV testing solely as a result of blood donation were considered as not having been tested for HIV. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.


DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, the percentage of U.S. adults who had ever been tested for HIV was 36.1% (95% confidence interval = 34.6%–37.6%), which was higher than, but not significantly different from, the 2004 estimate of 34.6%.

■ The annual percentage of adults who had ever had an HIV test increased from 30.2% in 1998 to 35.4% in 2003.



**Figure 10.2. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus, by age group and sex: United States, January–March 2005**




NOTES: The analyses excluded 311 adults (4.4%) with unknown human immunodeficiency virus (HIV) test status. Individuals who received HIV testing solely as a result of blood donation were considered as not having been tested for HIV.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who ever had an HIV test was highest among adults aged 25–34 years (56.1%) and lowest among adults aged 65 years and over (9.5%).
- For age groups 18–24 years, 25–34 years, and 35–44 years, women were more likely than men to have ever had an HIV test, whereas for age groups 45–64 years and 65 years and over, women were less likely than men to have ever had an HIV test.

**Figure 10.3. Age-sex-adjusted percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus, by race/ethnicity: United States, January–March 2005**



NOTES: The analyses excluded 311 adults (4.4%) with unknown human immunodeficiency virus (HIV) test status. Individuals who received HIV testing solely as a result of blood donation were considered as not having been tested for HIV. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The age-sex-adjusted percentages of persons who ever had an HIV test were 40.5% for Hispanic persons, 33.9% for non-Hispanic white persons, and 45.8% for non-Hispanic black persons.

- Of the three race/ethnicity groups, non-Hispanic black persons were most likely to have ever had an HIV test.

## Data tables for figures 10.1–10.3:

Data table for figure 10.1. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus: United States, 1997–2005

Year	Percent	95% confidence interval
1997	31.8	31.1-32.5
1998	30.2	29.5-30.9
1999	32.3	31.6-33.0
2000	32.1	31.4-32.8
2001	33.2	32.5-34.0
2002	34.5	33.8-35.2
2003	35.4	34.7-36.1
2004	34.6	33.9-35.3
January–March 2005	36.1	34.6-37.6

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 10.2. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus, by age group and sex: United States, January–March 2005**

<b>Age and sex</b>	<b>Percent</b>	<b>95% confidence interval</b>
<b>18–24 years</b>		
Total	34.3	30.0-38.7
Male	24.0	19.1-29.6
Female	44.7	38.6-50.9
<b>25–34 years</b>		
Total	56.1	52.6-59.5
Male	48.5	43.5-53.5
Female	63.5	58.9-67.9
<b>35–44 years</b>		
Total	48.8	45.7-52.0
Male	44.9	40.4-49.4
Female	52.7	48.5-56.8
<b>45–64 years</b>		
Total	30.7	28.5-33.0
Male	33.3	30.1-36.6
Female	28.3	25.3-31.5
<b>65 years and over</b>		
Total	9.5	7.7-11.5
Male	13.2	10.0-17.3
Female	6.6	4.8-9.0
<b>18 years and over: crude<sup>1</sup></b>		
Total	36.1	34.6-37.6
Male	34.4	32.3-36.5
Female	37.7	35.7-39.7
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	36.2	34.8-37.5
Male	34.0	32.1-35.9
Female	38.5	36.7-40.3

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 10.3. Age-sex-adjusted percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus, by race/ethnicity: United States, January–March 2005**

<b>Race/ethnicity</b>	<b>Percent<sup>1</sup></b>	<b>95% confidence interval</b>
<b>Hispanic or Latino</b>	40.5	37.1-43.9
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	33.9	32.3-35.6
<b>Black, single race</b>	45.8	42.1-49.5

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 11.1. Percentage of persons of all ages who assessed their health as excellent or very good: United States, 1997–2005**




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded persons with unknown health status (about 0.5% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, the percentage of persons who assessed their health as excellent or very good was 65.5% (95% confidence interval = 64.3%–66.7%), which was similar to the 2004 estimate of 66.5%.

■ From 1997 through 2001, the annual percentage of persons who assessed their health as excellent or very good remained similar: 68.5% in 1997, 69.1% in 1998, 68.7% in 1999, 68.3% in 2000, and 68.4% in 2001. The estimates decreased from 2001 to early 2005 (65.5%).

**Figure 11.2. Percentage distribution of respondent-assessed health status, by sex for all ages: United States, January–March 2005**




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 66 persons (0.31%) with unknown health status.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, most persons assessed their health as either excellent (35.0%) or very good (30.4%). A smaller percentage of persons assessed their health as good (25.0%). Fewer persons assessed their health as fair (7.1%) or poor (2.4%).

- Compared with males, females were more likely to assess their health as fair.

**Figure 11.3. Percentage of persons of all ages who assessed their health as excellent or very good, by age group and sex: United States, January–March 2005**




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 66 persons (0.31%) with unknown health status.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who assessed their health as excellent or very good decreased with age: 80.4% for those under 18 years, 65.0% for those aged 18–64 years, and 36.9% for those aged 65 years and over.
- There was no significant difference in the percentage of persons who assessed their health as excellent or very good between females and males for all three age groups.



**Figure 11.4. Age-sex-adjusted percentage of persons of all ages who assessed their health as excellent or very good, by race/ethnicity: United States, January–March 2005**



NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 66 persons (0.31%) with unknown health status. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ After adjusting for age and sex, the percentage of persons who assessed their health as excellent or very good was 56.1% for Hispanic persons, 69.4% for non-Hispanic white persons, and 56.6% for non-Hispanic black persons.

■ The health of Hispanic persons and non-Hispanic black persons was less likely to be assessed as excellent or very good than that of non-Hispanic white persons.

## Data tables for figures 11.1–11.4:

**Data table for figure 11.1. Percentage of persons of all ages who assessed their health as excellent or very good: United States, 1997–2005**

Year	Percent	95% confidence interval
1997	68.5	68.0-69.1
1998	69.1	68.6-69.6
1999	68.7	68.1-69.3
2000	68.3	67.7-68.8
2001	68.4	67.9-69.0
2002	67.4	66.8-68.0
2003	67.5	66.9-68.0
2004	66.5	66.0-67.1
January–March 2005	65.5	64.3-66.7

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 11.2. Percentage distribution of respondent-assessed health status, by sex for all ages: United States, January–March 2005**

Health status and sex	Percent	95% confidence interval
<b>Excellent</b>		
Total	35.0	33.8-36.3
Male	35.9	34.5-37.3
Female	34.2	32.8-35.7
<b>Very good</b>		
Total	30.4	29.4-31.5
Male	30.6	29.3-31.9
Female	30.3	29.2-31.5
<b>Good</b>		
Total	25.0	24.1-26.0
Male	24.7	23.6-25.8
Female	25.3	24.2-26.5
<b>Fair</b>		
Total	7.1	6.7-7.6
Male	6.6	6.1-7.2
Female	7.6	7.0-8.2
<b>Poor</b>		
Total	2.4	2.1-2.6
Male	2.2	1.9-2.6
Female	2.5	2.2-2.9

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 11.3. Percentage of persons of all ages who assessed their health as excellent or very good, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>Under 18 years</b>		
Total	80.4	78.6-82.0
Male	79.1	77.0-81.1
Female	81.6	79.7-83.5
<b>18–64 years</b>		
Total	65.0	63.6-66.3
Male	65.9	64.3-67.4
Female	64.1	62.5-65.6
<b>65 years and over</b>		
Total	36.9	34.5-39.3
Male	38.0	35.0-41.2
Female	36.0	33.1-39.0
<b>All ages: crude<sup>1</sup></b>		
Total	65.5	64.3-66.7
Male	66.5	65.2-67.8
Female	64.6	63.2-65.9
<b>All ages: age-adjusted<sup>2</sup></b>		
Total	65.4	64.3-66.5
Male	65.8	64.5-67.0
Female	65.1	63.8-66.3

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 11.4. Age-sex-adjusted percentage of persons of all ages who assessed their health as excellent or very good, by race/ethnicity: United States, January–March 2005**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
<b>Hispanic or Latino</b>	56.1	53.1-59.0
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	69.4	68.2-70.6
<b>Black, single race</b>	56.6	53.8-59.4

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997–2005**




NOTES: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home. The analyses excluded persons with unknown information on personal care needs (about 0.01% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the Appendix for more details.

DATA SOURCE: Family Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 7.0% (95% confidence interval = 6.0%–8.2%) of adults aged 65 years and over needed help with personal care from other persons. This estimate was higher than, but not significantly different from, the 2004 estimate of 6.3%.

■ The annual percentage of adults who needed help with personal care from other persons was 6.6% in 1997, 6.3% in 1998, 6.4% in 1999, 6.4% in 2000, 6.5% in 2001, 6.2% in 2002, 6.5% in 2003, and 6.3% in 2004.

**Figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January–March 2005**




NOTE: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, adults aged 85 years and over were more than six times as likely as adults aged 65–74 years old to need help with personal care from other persons (24.0% vs. 3.5%).
- There was no significant difference in the percentage of adults who needed help with personal care between women and men for all three age groups.

**Figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January–March 2005**



NOTES: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted percentage of persons who needed help with personal care from other persons was 11.2% for Hispanic persons, 6.4% for non-Hispanic white persons, and 9.1% for non-Hispanic black persons.

■ The percentage of those who needed help with personal care from other persons was higher for Hispanic persons than for non-Hispanic white persons.

## Data tables for figures 12.1–12.3:

Data table for figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997–2005

Year	Percent	95% confidence interval
1997	6.6	6.1-7.1
1998	6.3	5.9-6.8
1999	6.4	5.8-6.9
2000	6.4	5.9-6.9
2001	6.5	6.0-7.0
2002	6.2	5.7-6.7
2003	6.5	6.0-7.1
2004	6.3	5.8-6.8
January–March 2005	7.0	6.0-8.2

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January–March 2005

Age and sex	Percent	95% confidence interval
<b>65–74 years</b>		
Total	3.5	2.6-4.6
Male	2.9	1.8-4.7
Female	3.9	2.8-5.5
<b>75–84 years</b>		
Total	6.8	5.2-9.0
Male	4.8	3.0-7.6
Female	8.2	5.8-11.4
<b>85 years and over</b>		
Total	24.0	18.8-30.2
Male	18.7	11.9-28.2
Female	26.9	19.9-35.2
<b>65 years and over: crude<sup>1</sup></b>		
Total	7.0	6.0-8.2
Male	5.0	3.8-6.6
Female	8.5	7.0-10.3
<b>65 years and over: age-adjusted<sup>2</sup></b>		
Total	7.2	6.2-8.4
Male	5.5	4.2-7.2
Female	8.3	6.9-9.9

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January–March 2005**


<b>Race/ethnicity</b>	<b>Percent <sup>1</sup></b>	<b>95% confidence interval</b>
<b>Hispanic or Latino</b>	11.2	8.0-15.4
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	6.4	5.3-7.7
<b>Black, single race</b>	9.1	5.9-14.0

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



**Figure 13.1. Percentage of adults aged 18 years and over who experienced serious psychological distress during the past 30 days: United States, 1997–2005**




NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0–4) of the six items for each person are summed to yield a scale with a 0–to–24 range. A value of 13 or more for this scale is used here to define serious psychological distress (10). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 2.9% (95% confidence interval = 2.5%–3.4%) of adults aged 18 years and over experienced serious psychological distress during the past 30 days, which was similar to the 2004 estimate of 3.1%.

■ The annual percentage of adults who experienced serious psychological distress during the past 30 days declined significantly from 3.3% in 1997 to 2.4% in 1999, then increased from 2.4% in 1999 to 3.2% in 2001, and shows no trend in more recent years.

**Figure 13.2. Percentage of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by age group and sex: United States, January–March 2005**




NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0–4) of the six items for each person are summed to yield a scale with a 0–to–24 range. A value of 13 or more for this scale is used here to define serious psychological distress (10).

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, persons 65 years and over (1.8%) were less likely to have experienced serious psychological distress during the past 30 days compared with persons aged 18–44 years (2.9%) and 45–64 years (3.4%).
- There was no significant difference in the percentage of adults who experienced serious psychological distress during the past 30 days between women and men for all three age groups.

**Figure 13.3. Age-sex-adjusted percentage of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by race/ethnicity: United States, January–March 2005**



NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0–4) of the six items for each person are summed to yield a scale with a 0–to–24 range. A value of 13 or more for this scale is used here to define serious psychological distress (9). Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: 18–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted prevalence of serious psychological distress was 2.8% for Hispanic persons, 2.9% for non-Hispanic white persons, and 3.0% for non-Hispanic black persons.

■ The percentages of adults who experienced serious psychological distress during the past 30 days were similar across the three race/ethnicity groups.

## Data tables for figures 13.1–13.3:

**Data table for figure 13.1. Percentage of adults aged 18 years and over who experienced serious psychological distress during the past 30 days: United States, 1997–2005**

Year	Percent	95% confidence interval
1997	3.3	3.1-3.5
1998	3.0	2.7-3.2
1999	2.4	2.2-2.6
2000	2.7	2.5-2.9
2001	3.2	2.9-3.4
2002	3.0	2.8-3.2
2003	3.1	2.9-3.4
2004	3.1	2.9-3.4
January–March 2005	2.9	2.5-3.4

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 13.2. Percentage of adults aged 18 and over who experienced serious psychological distress during the past 30 days, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>18–44 years</b>		
Total	2.9	2.3-3.6
Male	2.5	1.7-3.6
Female	3.3	2.6-4.3
<b>45–64 years</b>		
Total	3.4	2.6-4.3
Male	3.0	2.0-4.5
Female	3.7	2.7-5.0
<b>65 years and over</b>		
Total	1.8	1.2-2.9
Male	-	-
Female	1.8	1.1-3.0
<b>18 years and over: crude<sup>1</sup></b>		
Total	2.9	2.5-3.4
Male	2.6	2.0-3.3
Female	3.2	2.6-3.8
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	2.9	2.4-3.3
Male	2.5	2.0-3.3
Female	3.2	2.6-3.8

-Estimates with a relative standard error greater than 30% are indicated with a dash, but data are not shown.

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 18–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 13.3. Age-sex-adjusted percentage of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by race/ethnicity: United States, January–March 2005**

<b>Race/ethnicity</b>	<b>Percent<sup>1</sup></b>	<b>95% confidence interval</b>
<b>Hispanic or Latino</b>	2.8	2.1-3.8
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	2.9	2.4-3.5
<b>Black, single race</b>	3.0	2.1-4.4

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–2005**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded persons with unknown diabetes status (about 0.1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 7.2% (95% confidence interval = 6.6%–8.0%) of U.S. adults aged 18 years and over had ever been diagnosed as having diabetes.

■ From 1997 to early 2005, there was an increasing trend in the prevalence of diagnosed diabetes among U.S. adults aged 18 years and over, from 5.1% in 1997 to 7.2% in early 2005.

**Figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, January–March 2005**




NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting “borderline” diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded 6 persons (0.1%) with unknown diabetes status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of diagnosed diabetes increased with age, with the highest rate among adults aged 65 years and over (16.5%) and the lowest rate among adults aged 18–44 years (2.1%).
- For the age group 55–64 years, the prevalence of diagnosed diabetes was higher among men than women.

**Figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, January–March 2005**



NOTES: Prevalence of diagnosed diabetes is based on self-report of ever having been diagnosed with diabetes by a doctor or other health professional. Persons reporting "borderline" diabetes status and women reporting diabetes only during pregnancy were not coded as having diabetes in the analyses. The analyses excluded six persons (0.1%) with unknown diabetes status. Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through March in the Sample Adult Core component of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted prevalence of diagnosed diabetes was 7.9% for Hispanic persons, 6.4% for non-Hispanic white persons, and 11.7% for non-Hispanic black persons.

■ The prevalence of diagnosed diabetes was higher among non-Hispanic black persons than Hispanic persons and non-Hispanic white persons.



## Data tables for figures 14.1–14.3:

**Data table for figure 14.1. Prevalence of diagnosed diabetes among adults aged 18 years and over: United States, 1997–2005**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	5.1 (4.9-5.4)	5.3 (5.1-5.6)
1998	5.3 (5.0-5.6)	5.4 (5.1-5.7)
1999	5.4 (5.1-5.7)	5.5 (5.2-5.8)
2000	5.9 (5.6-6.2)	6.0 (5.7-6.3)
2001	6.4 (6.1-6.7)	6.4 (6.1-6.7)
2002	6.5 (6.2-6.8)	6.5 (6.2-6.8)
2003	6.6 (6.3-6.9)	6.5 (6.2-6.9)
2004	7.0 (6.7-7.4)	6.9 (6.6-7.3)
January–March 2005	7.2 (6.6-8.0)	7.1 (6.5-7.8)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 14.2. Prevalence of diagnosed diabetes among adults aged 18 years and over, by age group and sex: United States, January–March 2005**

<b>Age and sex</b>	<b>Percent</b>	<b>95% confidence interval</b>
<b>18–44 years</b>		
Total	2.1	1.6-2.8
Male	2.3	1.5-3.4
Female	2.0	1.4-2.9
<b>45–54 years</b>		
Total	8.4	6.8-10.3
Male	10.4	8.1-13.4
Female	6.4	4.6-8.8
<b>55–64 years</b>		
Total	13.7	11.4-16.3
Male	15.9	12.4-20.2
Female	11.6	8.9-15.0
<b>65 years and over</b>		
Total	16.5	14.4-19.0
Male	17.7	14.4-21.6
Female	15.7	12.9-19.0
<b>18 years and over: crude<sup>1</sup></b>		
Total	7.2	6.6-8.0
Male	7.9	7.0-9.0
Female	6.6	5.8-7.6
<b>18 years and over: age-adjusted<sup>2</sup></b>		
Total	7.1	6.5-7.8
Male	8.0	7.0-9.0
Female	6.3	5.5-7.1

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.


**Data table for figure 14.3. Age-sex-adjusted prevalence of diagnosed diabetes among adults aged 18 years and over, by race/ethnicity: United States, January–March 2005**

<b>Race/ethnicity</b>	<b>Percent<sup>1</sup></b>	<b>95% confidence interval</b>
<b>Hispanic or Latino</b>	7.9	5.8-10.7
<b>Not Hispanic or Latino:</b>		
<b>White, single race</b>	6.4	5.8-7.2
<b>Black, single race</b>	11.7	9.3-14.6

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using four age groups: 18–44 years, 45–54 years, 55–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Figure 15.1. Percentage of persons of all ages who experienced an asthma episode in the past 12 months: United States, 1997–2005**




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded people with unknown asthma episode status (about 0.3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the Appendix for more details.

DATA SOURCE: Combined Sample Adult and Sample Child Core components of the 1997–2005 National Health Interview Surveys. The estimate for 2005 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, the percentage of persons of all ages who experienced an asthma episode in the past 12 months was 4.2% (95% confidence interval = 3.8%–4.8%), which was similar to the 2004 estimate of 4.1%.

■ The percentage of persons of all ages who experienced an asthma episode in the past 12 months decreased from 4.2% in 1997 to 3.9% in 1999, but increased significantly to 4.3% in 2001. The rates then decreased again to 3.9% in 2003.

**Figure 15.2. Percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and sex: United States, January–March 2005**




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 19 persons (0.2%) with unknown asthma episode status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who had an asthma episode in the past 12 months was higher among children under age 15 years than among those aged 15 years and over. This age difference was also seen among males, but not among females.
- For children under age 15 years, the percentage of persons who had an asthma episode in the past 12 months was higher among boys than among girls. However, for persons 15 years and over, the percentage was higher among women than among men.

**Figure 15.3. Sex-adjusted percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and race/ethnicity: United States, January–March 2005**




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 19 persons (0.2%) with unknown asthma episode status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For children under age 15 years, the sex-adjusted percentage of persons who had an asthma episode in the past 12 months was higher among non-Hispanic black children than among Hispanic or non-Hispanic white children, but this difference was not statistically significant.
- Among the six age-specific race/ethnicity groups, the sex-adjusted percentage of persons who had an asthma episode in the past 12 months was highest among non-Hispanic black children under age 15 years (7.5%).

**Figure 15.4. Prevalence of current asthma among persons of all ages: United States, 2001–2004**




NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded people with unknown current asthma status (about 0.3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2001–02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the Appendix for more details.

DATA SOURCE: Combined Sample Adult and Sample Child Core components of the 2001–2005 National Health Interview Surveys. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2005, 7.6% (95% confidence interval = 7.0%–8.2%) of persons of all ages currently had asthma, which was higher than, but not significantly different from, the 2004 estimate of 7.3%.

■ The prevalence of current asthma among persons of all ages decreased from 7.6% in 2001 to 7.1% in 2003.

**Figure 15.5. Prevalence of current asthma among persons of all ages, by age group and sex: United States, January–March 2005**




NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 21 persons (0.2%) with unknown current asthma status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of current asthma was higher among children 0–14 years old than among persons 15 years and over. This pattern was seen among males but not among females.
- For children under age 15 years, the prevalence of current asthma was higher among boys than among girls. However, among those 15 years and over, the prevalence of current asthma was higher among women than among men.

**Figure 15.6. Sex-adjusted prevalence of current asthma among persons of all ages, by age group and race/ethnicity: United States, January–March 2005**



NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 21 persons (0.2%) with unknown current asthma status.

DATA SOURCE: Based on data collected from January through March in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For children under age 15 years, the sex-adjusted prevalence of current asthma was higher among non-Hispanic black children than among Hispanic or non-Hispanic white children.

■ Among the six age-specific race/ethnicity groups, the sex-adjusted prevalence of current asthma was highest among non-Hispanic black children under age 15 years (15.7%).



## Data tables for figures 15.1–15.6:

**Data for figure 15.1. Percentage of persons of all ages who experienced an asthma episode in the past 12 months: United States, 1997–2005**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
1997	4.2 (4.0-4.4)	4.2 (3.9-4.4)
1998	4.0 (3.7-4.2)	3.9 (3.7-4.2)
1999	3.9 (3.6-4.1)	3.9 (3.6-4.1)
2000	4.0 (3.8-4.2)	4.0 (3.8-4.2)
2001	4.3 (4.1-4.6)	4.3 (4.1-4.5)
2002	4.3 (4.0-4.5)	4.3 (4.0-4.5)
2003	3.9 (3.7-4.1)	3.9 (3.7-4.1)
2004	4.1 (3.8-4.3)	4.1 (3.8-4.3)
January–March 2005	4.2 (3.8-4.8)	4.2 (3.8-4.8)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 15.2. Percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>0–14 years</b>		
Total	5.8	4.7-7.2
Male	7.1	5.6-9.1
Female	4.5	3.2-6.2
<b>15–34 years</b>		
Total	3.9	3.0-5.0
Male	2.8	1.8-4.3
Female	5.0	3.6-6.8
<b>35 years and over</b>		
Total	3.8	3.2-4.5
Male	2.9	2.2-3.9
Female	4.5	3.7-5.5
<b>All ages: crude<sup>1</sup></b>		
Total	4.2	3.8-4.8
Male	3.8	3.2-4.5
Female	4.6	4.0-5.4
<b>All ages: age-adjusted<sup>2</sup></b>		
Total	4.2	3.8-4.8
Male	3.8	3.2-4.5
Female	4.6	3.9-5.4

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 15.3. Sex-adjusted percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and race/ethnicity: United States, January–March 2005**

Age and race/ethnicity	Percent <sup>1</sup>	95% confidence interval
<b>0–14 years</b>		
Hispanic or Latino	5.9	3.9-8.8
Not Hispanic or Latino:		
White, single race	5.1	3.8-6.7
Black, single race	7.5	4.4-12.4
<b>15 years and over</b>		
Hispanic or Latino	3.0	2.0-4.3
Not Hispanic or Latino:		
White, single race	3.8	3.3-4.5
Black, single race	3.7	2.6-5.3

<sup>1</sup>Estimates are sex-adjusted to the 2000 projected U.S. standard population.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data for figure 15.4. Prevalence of current asthma among persons of all ages: United States, 2001–2004**

Year	Percent (95% confidence interval)	
	Crude <sup>1</sup>	Age-adjusted <sup>2</sup>
2001	7.6 (7.3-7.9)	7.6 (7.3-7.9)
2002	7.4 (7.1-7.7)	7.4 (7.1-7.7)
2003	7.1 (6.8-7.4)	7.1 (6.8-7.4)
2004	7.3 (7.0-7.6)	7.3 (7.1-7.7)
January–March 2005	7.6 (7.0-8.2)	7.6 (7.0-8.2)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 2001–05. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 15.5. Prevalence of current asthma among persons of all ages, by age group and sex: United States, January–March 2005**

Age and sex	Percent	95% confidence interval
<b>0–14 years</b>		
Total	9.3	8.0-10.9
Male	10.8	8.8-13.1
Female	7.8	6.1-9.9
<b>15–34 years</b>		
Total	7.0	5.9-8.4
Male	5.1	3.7-6.8
Female	9.0	7.1-11.2
<b>35 years and over</b>		
Total	7.2	6.4-8.1
Male	5.5	4.5-6.7
Female	8.7	7.6-10.1
<b>All ages: crude<sup>1</sup></b>		
Total	7.6	7.0-8.2
Male	6.5	5.8-7.4
Female	8.6	7.7-9.6
<b>All ages: age-adjusted<sup>2</sup></b>		
Total	7.6	7.0-8.2
Male	6.5	5.7-7.3
Female	8.6	7.7-9.6

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for figure 15.6. Sex-adjusted prevalence of current asthma among persons of all ages, by age group and race/ethnicity: United States, January–March 2005**

Age and race/ethnicity	Percent <sup>1</sup>	95% confidence interval
<b>0–14 years</b>		
Hispanic or Latino	10.1	7.4-13.6
Not Hispanic or Latino:		
White, single race	6.9	5.5-8.7
Black, single race	15.7	11.5-21.0
<b>15 years and over</b>		
Hispanic or Latino	5.5	4.2-7.3
Not Hispanic or Latino:		
White, single race	6.9	6.2-7.7
Black, single race	8.9	6.9-11.4

<sup>1</sup>Estimates are sex-adjusted to the 2000 projected U.S. standard population.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

## References

1. U.S. Department of Health and Human Services. Healthy People 2010: Understanding and improving health. Washington: U.S. Department of Health and Human Services. Washington. 2000.
2. Office of Management and Budget. Revisions to the standards for the classification of Federal data on race/ethnicity. Executive Office of the President, Office of Management and Budget (OMB), Office of Information and Regulatory Affairs. Available at <http://www.whitehouse.gov/omb/fedreg>. Published on October 30, 1997 and accessed on September 11, 2003.
3. National Center for Health Statistics. Health, United States, 2003. Hyattsville, MD: 2003.
4. Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. Healthy People Statistical Notes, no 20. Hyattsville, MD. National Center for Health Statistics. 2001.
5. National Center for Health Statistics (assorted authors). Early release of selected estimates based on data from the National Health Interview Survey. National Center for Health Statistics. Available at: <http://www.cdc.gov/nchs/nhis.htm>. Series of Internet-only reports released approximately quarterly beginning in April 2001.
6. Centers for Disease Control and Prevention. Interim Influenza Vaccination Recommendations, 2004--05 Influenza Season. MMWR 2004; 53(39):923-924.
7. Centers for Disease Control and Prevention. Notice to readers: Updated recommendations from the advisory committee on immunization practices in response to delays in supply of influenza vaccine for the 2000-01 season. MMWR 2000; 49(39):888-92.
8. Centers for Disease Control and Prevention. Notice to readers: Delayed influenza vaccine availability for 2001-02 season and supplemental recommendations of the advisory committee on immunization practices. MMWR 2001; 50(27):582-5.
9. Centers for Disease Control and Prevention. Prevention and control of influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 2003; 52(RR08):1-36
10. Kessler RC, Barker PR, Colpe LJ, et al. Screening for serious mental illness in the general population. Arch Gen Psychiatry. 60:184-9. 2003.

**Table I. The 2005 National Health Interview Survey questions used to define selected health measures**

**Lack of health insurance and type of coverage<sup>1</sup>**

FHI.050<sup>2</sup> **Field Representative: SHOW FLASHCARD F10**

{Are you/Is anyone} covered by any kind of health insurance or some other kind of health care plan?

**Field Representative: READ IF NECESSARY: INCLUDE HEALTH INSURANCE OBTAINED THROUGH EMPLOYMENT OR PURCHASED DIRECTLY AS WELL AS GOVERNMENT PROGRAMS LIKE MEDICARE AND MEDICAID THAT PROVIDE MEDICAL CARE OR HELP PAY MEDICAL BILLS.**

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

FHI.070 What kind of health insurance or health care coverage {do/does} {you/subject name} have? **INCLUDE** those that pay for only one type of service (nursing home care, accidents, or dental care), **exclude** private plans that only provide extra cash while hospitalized.

**Field Representative: PLEASE REFER TO FLASHCARDS F12 AND F13 FOR YOUR STATE.**

- (01) Private health insurance
- (02) Medicare
- (03) Medi-Gap
- (04) Medicaid
- (05) SCHIP (CHIP/Children's Health Insurance Program)
- (06) Military health care (TRICARE/VA/CHAMP-VA)
- (07) Indian Health Service
- (08) State-sponsored health plan
- (09) Other government program
- (10) Single service plan (e.g. dental, vision, prescriptions)
- (11) No coverage of any type
- (97) Refused
- (99) Don't know

FHI.072 People covered by Medicare have a card that looks like this. {Are/Is} {person} covered by Medicare?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

- FHI.073 There is a program called Medicaid that pays for health care for persons in need. In this state it is also called [state name]. {Are/Is} {person} covered by Medicaid?
- (1) Yes
  - (2) No
  - (7) Refused
  - (9) Don't know

**Usual place to go for medical care**

- AAU.020 Is there a place that you USUALLY go to when you are sick or need advice about your health?

- (1) Yes
- (2) There is NO place
- (3) There is MORE THAN ONE place
- (7) Refused
- (9) Don't know

- AAU.030 What kind of place is it—a clinic, doctor's office, emergency room, or some other place?

- (1) Clinic or health center
- (2) Doctor's office or HMO
- (3) Hospital emergency room
- (4) Hospital outpatient department
- (5) Some other place
- (6) Doesn't go to one place most often
- (7) Refused
- (9) Don't know

**Obtaining needed medical care**

- FAU.040 DURING THE PAST 12 MONTHS, was there any time when {you/someone in the family} needed medical care, but did not get it because {you/the family} couldn't afford it?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

**Influenza vaccination**

- AAU.310 DURING THE PAST 12 MONTHS, have you had a flu shot? A flu shot is usually given in the fall and protects against influenza for the flu season.

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

AAU.315 DURING THE PAST 12 MONTHS, have you had a flu vaccine sprayed in your nose by a doctor or other health professional? A health professional may have let you spray it. This vaccine is usually given in the fall and protects against influenza for the flu season.

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

**Pneumococcal vaccination**

AAU.320 Have you EVER had a pneumonia shot? This shot is usually given only once or twice in a person's lifetime and is different from the flu shot. It is also called the pneumococcal vaccine.

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

**Obesity**

AHB.190 How tall are you without shoes?

- (2-7) 2-7 feet
- (0-11) 0-11 Inches
- (97) Refused
- (99) Don't know

FT Feet \_\_\_\_\_  
IN Inches \_\_\_\_\_

AHB.200 How much do you weigh without shoes?

- (50-500) 50-500 pounds
- (997) Refused
- (999) Don't know

### Leisure-time physical activity

The next questions are about physical activities (exercises, sports, physically active hobbies...) that you may do in your LEISURE time.

AHB.110 How often do you do LIGHT OR MODERATE activities for AT LEAST 10 MINUTES that cause ONLY LIGHT sweating or a SLIGHT to MODERATE increase in breathing or heart rate?

**Field Representative: IF NECESSARY, PROMPT WITH: HOW MANY TIMES PER DAY, PER WEEK, PER MONTH, OR PER YEAR DO YOU DO THESE ACTIVITIES?**

@NO times per

(0) Never

(1-995) 1-995

(996) Unable to do this type activity

(997) Refused

(999) Don't know

@TP time period

(1) Day

(2) Week

(3) Month

(4) Year

AHB.120 About how long do you do these light or moderate activities each time?

@NO number

(1-995) 1-995

(997) Refused

(999) Don't know

@TP time period

(1) Minutes

(2) Hours

(7) Refused

(9) Don't know



AHB.090 How often do you do VIGOROUS activities for AT LEAST 10 MINUTES that cause HEAVY sweating or LARGE increases in breathing or heart rate?

**Field Representative: IF NECESSARY, PROMPT WITH: HOW MANY TIMES PER DAY, PER WEEK, PER MONTH, OR PER YEAR DO YOU DO THESE ACTIVITIES?**

@NO times per

(0) Never

(1-995) 1-995

(996) Unable to do this type activity

(997) Refused

(999) Don't know

@TP time period

(1) Day

(2) Week

(3) Month

(4) Year

AHB.100 About how long do you do these vigorous activities each time?

@NO number

(1-995) 1-995

(997) Refused

(999) Don't know

@TP time period

(1) Minutes

(2) Hours

(7) Refused

(9) Don't know

**Current smoking**

AHB.010 Have you smoked at least 100 cigarettes in your ENTIRE LIFE?

(1) Yes

(2) No

(7) Refused

(9) Don't know

AHB.030 Do you NOW smoke cigarettes every day, some days or not at all?

(1) Every day

(2) Some days

(3) Not at all

(7) Refused

(9) Don't know

**Alcohol consumption**

AHB.150 In your ENTIRE LIFE, have you had at least 12 drinks of any type of alcoholic beverage?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

AHB.180 In the PAST YEAR, on how many DAYS did you have 5 or more drinks of any alcoholic beverage?

- (0) Never
- (1-365) 1-365
- (997) Refused
- (999) Don't know

**HIV testing**

ADS.040 Except for tests you may have had as part of blood donations, have you ever been tested for HIV?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

**General health status**

FHS.500 Would you say {subject name's} health in general is excellent, very good, good, fair, or poor?

- (1) Excellent
- (2) Very good
- (3) Good
- (4) Fair
- (5) Poor
- (7) Refused
- (9) Don't know

**Personal care needs**

FHS.070 Because of a physical, mental, or emotional problem, {do/does} {you/anyone in the family} need the help of other persons with PERSONAL CARE NEEDS, such as eating, bathing, dressing, or getting around inside this home?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

**Serious psychological distress**

ACN.471 During the PAST 30 DAYS, how often did you feel...

- ...So sad that nothing could cheer you up?
- ...Nervous?
- ...Restless or fidgety?
- ...Hopeless?
- ...That everything was an effort?
- ...Worthless?

- (1) ALL of the time
- (2) MOST of the time
- (3) SOME of the time
- (4) A LITTLE of the time
- (5) NONE of the time
- (7) Refused
- (9) Don't know

**Diagnosed diabetes**

ACN.160 Other than during pregnancy, have you EVER been told by a doctor or health professional that you have diabetes or sugar diabetes?

- (1) Yes
- (2) No
- (3) Borderline
- (7) Refused
- (9) Don't know

**Asthma**

ACN.090 DURING THE PAST 12 MONTHS, have you had an episode of asthma or an asthma attack?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

ACN.085 Do you still have asthma?

- (1) Yes
- (2) No
- (7) Refused
- (9) Don't know

<sup>1</sup>Information from follow-up questions such as plan name(s) was used to reassign insurance status and type of coverage to avoid misclassification.

<sup>2</sup>Alphanumeric codes refer directly to the question on the 2005 NHIS that was used to define the health measure.

**Table II. Number and percentage of uninsured persons, persons with public coverage, and persons with private coverage, based on the 1990 and 2000 census population estimates, by age group: United States, 2000–02**

Health insurance coverage and age group	2000		2001		2002	
	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>	1990-based <sup>1</sup>	2000-based <sup>2</sup>
<b>Number of uninsured in millions</b>						
All ages	40.4	41.3	39.1	40.2	40.4	41.5
Under 65 years	39.9	40.8	38.7	39.8	40.0	41.1
18–64 years	31.2	32.0	30.9	31.9	32.4	33.5
Under 18 years	8.8	8.9	7.8	7.9	7.6	7.6
<b>Percent (95% confidence interval)</b>						
<b>Uninsured</b>						
All ages	14.7 (14.3-15.1)	14.9 (14.5-15.3)	14.1 (13.7-14.6)	14.3 (13.8-14.8)	14.5 (14.1-14.9)	14.7 (14.3-15.1)
Under 65 years						
Crude	16.6 (16.1-17.1)	16.8 (16.3-17.2)	15.9 (15.4-16.4)	16.2 (15.7-16.7)	16.3 (15.8-16.7)	16.5 (16.0-16.9)
Age-adjusted	16.6 (16.1-17.1)	16.8 (16.3-17.3)	16.0 (15.5-16.5)	16.2 (15.7-16.7)	16.4 (15.9-16.9)	16.6 (16.1-17.1)
18–64 years	18.4 (17.9-18.9)	18.7 (18.1-19.2)	18.1 (17.5-18.6)	18.3 (17.8-18.8)	18.8 (18.3-19.3)	19.1 (18.6-19.6)
Under 18 years	12.2 (11.6-12.8)	12.3 (11.7-12.9)	10.8 (10.1-11.5)	11.0 (10.3-11.7)	10.4 (9.8-11.0)	10.5 (9.9-11.1)
<b>Public</b>						
Under 65 years	12.9 (12.4-13.4)	12.9 (12.4-13.4)	13.6 (13.1-14.1)	13.6 (13.1-14.1)	15.2 (14.6-15.8)	15.2 (14.6-15.8)
18–64 years	9.0 (8.7-9.4)	9.1 (8.7-9.4)	9.4 (9.0-9.8)	9.4 (9.0-9.8)	10.2 (9.8-10.7)	10.3 (9.9-10.7)
Under 18 years	21.8 (20.8-22.8)	22.0 (21.0-23.0)	23.4 (22.4-24.3)	23.6 (22.6-24.5)	26.8 (25.7-27.9)	27.1 (26.0-28.2)
<b>Private</b>						
Under 65 years	72.0 (71.4-72.7)	71.8 (71.1-72.5)	71.9 (71.2-72.6)	71.6 (70.9-72.3)	70.1 (69.3-70.9)	69.8 (69.0-70.6)
18–64 years	74.0 (73.4-74.6)	73.8 (73.2-74.4)	73.9 (73.3-74.6)	73.7 (73.1-74.4)	72.6 (71.9-73.2)	72.3 (71.6-72.9)
Under 18 years	67.5 (66.4-68.5)	67.1 (66.1-68.2)	67.1 (66.0-68.2)	66.7 (66.4-68.6)	64.3 (63.1-65.5)	63.9 (62.7-65.1)

<sup>1</sup>The estimates are calculated based on data with sample weights derived from the 1990 census.

<sup>2</sup>The estimates are calculated based on data with sample weights derived from the 2000 census.

DATA SOURCE: National Health Interview Survey, 2000–02. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Table III. Selected estimates using National Health Interview Survey weights based on the 1990 and 2000 census population estimates: United States, 2000–02**

Indicator	2000		2001		2002	
	1990 based <sup>1</sup>	2000 based <sup>2</sup>	1990 based <sup>1</sup>	2000 based <sup>2</sup>	1990 based <sup>1</sup>	2000 based <sup>2</sup>
<b>Percent (95% confidence interval)</b>						
With a usual place to go for medical care	87.1 (86.7-87.5)	87.0 (86.5-87.4)	88.2 (87.7-88.6)	88.0 (87.6-88.5)	87.8 (87.3-88.2)	87.6 (87.1-88.0)
Failed to obtain needed care due to cost	4.5 (4.3-4.7)	4.5 (4.3-4.7)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)
Had influenza shot in the past year	64.3 (62.9-65.7)	64.4 (63.0-65.9)	63.0 (61.6-64.4)	63.1 (61.7-64.5)	65.6 (64.1-67.0)	65.7 (64.3-67.2)
Had pneumococcal vaccination in the past year	52.9 (51.3-54.5)	53.1 (51.5-54.7)	53.8 (52.2-55.4)	54.0 (52.4-55.6)	55.7 (54.2-57.2)	56.0 (54.5-57.5)
With obesity	21.8 (21.2-22.4)	21.8 (21.2-22.4)	23.0 (22.4-23.5)	23.0 (22.4-23.6)	23.9 (23.3-24.6)	23.9 (23.3-24.6)
Engaged in regular leisure-time physical activity	32.0 (31.3-32.7)	31.9 (31.2-32.6)	32.1 (31.3-32.8)	32.1 (31.3-32.8)	32.0 (31.2-32.8)	31.9 (31.1-32.7)
Current smoking	23.3 (22.7-23.9)	23.2 (22.5-23.8)	22.8 (22.2-23.4)	22.7 (22.1-23.3)	22.5 (21.9-23.1)	22.4 (21.7-23.0)
Had 5 or more drinks in one day at least once in the past year	19.3 (18.7-20.0)	19.2 (18.6-19.9)	20.1 (19.5-20.7)	20.0 (19.4-20.6)	19.9 (19.3-20.6)	19.9 (19.2-20.5)
Ever had HIV <sup>3</sup> test	32.2 (31.5-32.9)	32.1 (31.4-32.8)	33.3 (32.6-34.0)	33.2 (32.5-34.0)	34.5 (33.8-35.3)	34.5 (33.8-35.2)
With excellent or very good health status	68.4 (67.9-69.0)	68.3 (67.7-68.8)	68.6 (68.1-69.2)	68.4 (67.9-69.0)	67.6 (67.0-68.2)	67.4 (66.8-68.0)
With need for personal care	6.3 (5.9-6.8)	6.4 (5.9-6.9)	6.5 (6.0-7.0)	6.5 (6.0-7.0)	6.2 (5.6-6.7)	6.2 (5.7-6.7)
Serious psychological distress	2.7 (2.5-2.9)	2.7 (2.5-2.9)	3.2 (2.9-3.4)	3.2 (2.9-3.4)	3.0 (2.8-3.3)	3.0 (2.8-3.2)
Diagnosed diabetes	5.9 (5.6-6.2)	5.9 (5.6-6.2)	6.4 (6.1-6.7)	6.4 (6.1-6.7)	6.5 (6.2-6.8)	6.5 (6.2-6.8)
Asthma episodes in the past year	40.1 (37.9-42.4)	40.0 (37.7-42.2)	43.4 (41.0-45.8)	43.2 (40.8-45.5)	42.8 (40.4-45.2)	42.6 (40.2-44.9)

<sup>1</sup>The estimates are calculated based on data with sample weights derived from the 1990 census.

<sup>2</sup>The estimates are calculated based on data with sample weights derived from the 2000 census.

<sup>3</sup>HIV is human immunodeficiency virus.

DATA SOURCE: National Health Interview Survey, 2000–02. Data are based on household interviews of a sample of the civilian noninstitutionalized population.